

Quantum Antenna Hypothesis

M. Pitkänen¹, February 1, 2006

¹ Department of Physical Sciences, High Energy Physics Division,
PL 64, FIN-00014, University of Helsinki, Finland.
matpitka@rock.helsinki.fi, <http://www.physics.helsinki.fi/~matpitka/>.
Recent address: Puutarhurinkatu 10,10960, Hanko, Finland.

Contents

| | | |
|----------|--|-----------|
| 1 | Introduction | 4 |
| 1.1 | Massless extremals and quantum antenna hypothesis | 4 |
| 1.2 | Evidence | 5 |
| 1.3 | Quantum antenna hypothesis and brain consciousness | 6 |
| 1.3.1 | Brain as a neuron society metaphor | 6 |
| 1.3.2 | The notion of neural window | 6 |
| 1.3.3 | Music metaphor | 7 |
| 1.3.4 | Brain as an associative net | 7 |
| 1.4 | Relationship of TGD approach with microtubular approach | 7 |
| 1.5 | MEs and information molecules | 8 |
| 1.6 | MEs and quantum holography | 8 |
| 1.7 | MEs and the notion of conscious hologram | 9 |
| 1.8 | Negative energy MEs and bio-control | 10 |
| 1.9 | MEs and dark matter hierarchy | 10 |
| 2 | Massless extremals | 11 |
| 2.1 | Massless extremals as general solutions of field equations | 11 |
| 2.2 | About the electro-weak and color fields associated with massless extremals | 12 |
| 2.3 | How massless extremals generate coherent states of photons? | 13 |
| 2.4 | Massless extremal is accompanied by a Bose-Einstein condensate of parallel photons | 14 |
| 2.5 | MEs and exotic representations of p-adic Super Virasoro | 15 |
| 3 | Microtubules as quantum antennae | 16 |
| 3.1 | Linear structures as quantum antennae | 16 |
| 3.2 | Are microtubules accompanied by massless extremals? | 16 |
| 3.3 | How macroscopic quantum coherence is generated? | 18 |
| 3.3.1 | Topological field quantization | 18 |
| 3.3.2 | Phase locking for the system of Josephson junctions | 18 |
| 3.3.3 | Gap junctions and MAPs | 19 |
| 3.4 | Are nerve pulse patterns coded into vacuum currents and coherent light? | 19 |
| 4 | Massless extremals and information molecules | 20 |
| 4.1 | Questions about information molecules | 20 |
| 4.2 | A model of biological self-organization based on quantum antenna hypothesis | 21 |

| | | |
|----------|--|-----------|
| 5 | Evidence for quantum antenna hypothesis | 22 |
| 5.1 | TGD inspired model for sonoluminescence | 22 |
| 5.1.1 | Sonoluminescence and hydrodynamic hierarchy | 23 |
| 5.1.2 | The model of sonoluminescence by Buzzacchi, del Giudice and Preparata | 24 |
| 5.1.3 | TGD inspired model for sonoluminescence | 25 |
| 5.2 | Stirred and shaken | 28 |
| 5.3 | Evidence for quantum antenna hypothesis in living systems | 29 |
| 5.4 | Biefeld-Brown effect | 30 |
| 6 | Quantum holography and MEs | 32 |
| 6.1 | Quantum holography in the sense of quantum gravity theories | 33 |
| 6.2 | Generalization of the solution ansatz defining massless extremals (MEs) | 34 |
| 6.2.1 | Local light cone coordinates | 34 |
| 6.2.2 | A conformally invariant family of local light cone coordinates | 35 |
| 6.2.3 | Closer inspection of the conditions defining local light cone coordinates | 35 |
| 6.2.4 | General solution ansatz for MEs for given choice of local light cone coordinates | 36 |
| 6.3 | Quantum holography in non-cosmological length scales | 37 |
| 6.4 | Supercanonical representations, quantum holography, and consciousness | 38 |
| 6.5 | Quantum holography and quantum information theory | 38 |
| 6.6 | MEs and qualia | 40 |
| 6.7 | Connection with the notion of wave-DNA | 40 |
| 6.7.1 | The notion of wave-DNA | 41 |
| 6.7.2 | DNA MEs as counterparts of wave DNA | 41 |
| 6.7.3 | DNA solitons and phantom DNA effect as evidence for superconducting DNA space-time sheets | 41 |
| 6.8 | p-Adic cognition and the existence of double-sheeted MEs and wormhole magnetic fields | 42 |
| 6.9 | Connection with quantum holography involved with NMR | 43 |
| 6.10 | Summary | 44 |
| 7 | Appendix: A model for the topological condensation of coherent vapor phase photons | 44 |
| 7.1 | The action | 45 |
| 7.2 | Coherent state is generated in resonant-like manner for light-like vacuum currents | 46 |
| 7.3 | Stimulated topological condensation | 47 |

Abstract

So called massless extremals are nonvacuum extremals of both Kähler action and the EYM action serving as effective action of the theory. These extremals have cylindrical geometry and are carriers of purely classical vacuum currents and Einstein tensor, which are both light like. These vacuum currents generate coherent states of photons and gravitons with frequencies coming as multiples of the basic frequency determined by the length of the microtubule. It is proposed that microtubules and other linear structures could act as quantum antennae so that coherent light is for brain same as radiowaves for us. Massless extremals associated with axonal microtubules or axons themselves could serve as waveguides for the photons of coherent light and realize the notion of neural window abstracted from the paradigm of holographic brain. Vacuum currents could be also behind the ability of the biosystems to form representations of the external world.

There is indeed evidence for the quantum antenna hypothesis: some monocellulars are known to possess primitive microtubular vision, biophotons of Popp could be generated by massless extremals and the observations of Callahan support the view that odour perception of insects relies on maserlike emissions by the odour molecules. The coherent light emitted in sonoluminescence could be generated by lightlike vacuum currents associated with regions with size given roughly by the diameter of microtubule when vapour-to-liquid phase transition occurs at the final stage of the bubble collapse. Also the observed direct transformation of kinetic energy of fluid motion to chemical energy could involve generation of massless extremals.

The lightlike boundaries of MEs have the same miraculous conformal properties as the boundary of future lightcone and MEs also allow holography in the sense of quantum gravity and string models and there are good hopes to generalize the construction of the configuration space geometry and quantum TGD to take into account the classical non-determinism of Kähler action. MEs provide a justification for the intuition that the supercanonical and superconformal symmetries of the lightcone boundary $\delta M_+^4 \times CP_2$, which are cosmological symmetries, generalize to approximate macroscopic symmetries acting on the lightlike boundaries of the spacetime sheets inside future lightcone and broken only by quantum gravity. Supercanonical symmetries almost-commute with Poincare symmetries and the gigantic almost-degenerate supercanonical multiplets defined by genuinely quantum gravitational state functionals in the 'world of worlds' correspond in a well-defined sense to higher abstraction level expected to be crucial for understanding consciousness. MEs are also tailor-made for quantum holography and teleportation. Quantum holography conceptualization inspires much more detailed views about how biosystems process information and how this information becomes conscious.

So called massless extremals are nonvacuum extremals of both Kähler action and the EYM action serving as effective action of the theory. These extremals have cylindrical geometry and are carriers of purely classical vacuum currents and Einstein tensor, which are both light like. These vacuum currents generate coherent states of photons and gravitons with frequencies coming as multiples of the basic frequency determined by the length of the microtubule. It is proposed that microtubules and other linear structures could act as quantum antennae so that coherent light is for brain same as radiowaves for us. Massless extremals associated with axonal microtubules or axons themselves could serve as waveguides for the photons of coherent light and realize the notion of neural window abstracted from the paradigm of holographic brain. Vacuum currents could be also behind the ability of the biosystems to form representations of the external world.

There is indeed evidence for the quantum antenna hypothesis: some monocellulars are known to possess primitive microtubular vision, biophotons of Popp could be generated by massless extremals and the observations of Callahan support the view that odour perception of insects relies on maserlike emissions by the odour molecules. The coherent light emitted in sonoluminescence could be generated by lightlike vacuum currents associated with regions with size given roughly by the diameter of microtubule when vapour-to-liquid phase transition occurs at the final stage of the bubble collapse. Also the observed direct transformation of kinetic energy of fluid motion to chemical energy could involve generation of massless extremals.

The lightlike boundaries of MEs have the same miraculous conformal properties as the boundary of future lightcone and MEs also allow holography in the sense of quantum gravity and string models and there are good hopes to generalize the construction of the configuration space geometry and quantum TGD to take into account the classical non-determinism of Kähler action. MEs provide a justification for the intuition that the supercanonical and superconformal symmetries of the lightcone boundary $\delta M_+^4 \times CP_2$, which are cosmological symmetries, generalize to approximate macroscopic symmetries acting on the lightlike boundaries of the spacetime sheets inside future lightcone and broken only by quantum gravity. Supercanonical symmetries almost-commute with Poincare symmetries and the gigantic almost-degenerate supercanonical multiplets defined by genuinely quantum gravitational state functionals in the 'world of worlds' correspond in a well-defined sense to higher abstraction level expected to be crucial for understanding consciousness. MEs are also tailor-made for quantum holography and teleportation. Quantum holography conceptualization inspires much more detailed views about how biosystems process information and how this information becomes conscious.

1 Introduction

One of the basic problems faced by the quantum theories of consciousness is to understand how macroscopic quantum coherence in the brain is realized. Bose-Einstein condensates and coherent states are believed to be crucial in this respect but the great problem is how macroscopic quantum phases could be realized in the wetty, noisy and hot environment provided by brain. In TGD framework the notion of many-sheeted space-time provides a solution to this basic problem. Furthermore, the notion of self as a subsystem able to remain p-adically un-entangled is consistent with the fact that macroscopic quantum phases behave like quantum particles. The general views about macroscopic quantum phases predicted by TGD and about their role with regards to consciousness is described in previous chapters. This chapter is devoted to coherent and Bose-Einstein condensed photons which are crucial in the quantum models of brain consciousness relating on microtubules and seem to be associated with linear structures also in TGD framework. These linear structures include not only microtubules but also axons, DNA and proteins and most of the considerations to follow are quite general and by no means restricted to microtubules.

1.1 Massless extremals and quantum antenna hypothesis

The purpose of this chapter is a more detailed formulation of the quantum antenna hypothesis stating that microtubules generate a macroscopic coherent state of photons. The so called massless extremals are a very general class of zero action non-vacuum extremals of both the Kähler action and the effective action and differentiate clearly between TGD and standard gauge theories, in particular QED. Massless extremals describe the propagation of massless gauge fields in single preferred direction. The polarization for given values of transversal coordinates has a fixed direction. Linear superposition is not possible.

Topological field quantization assigns to each quantum notion its classical counterpart and a very attractive identification of the massless extremals is as the classical counterparts of massless classical quanta such as photons and gravitons. Even the classical counterparts of the virtual particles make sense: in particular, negative energy photons represented by annihilation part of the free photon field seem to have geometric representation as negative energy massless extremals.

Massless extremals (ME) of the effective action can indeed generate coherent states of photons and gravitons.

1. ME:s are characterised by light like vacuum Kähler current J_K (this is purely TGD:eish feature!). In general but not always this implies light-like em current J_{em} and the standard coupling to the quantized photon field generates a coherent state of photons.

2. The geometry of the 3-surface in question is most naturally cylindrical and microtubules (as also DNA, proteins, etc..) indeed possess this kind of geometry. There are sharp resonances at frequencies $\omega = n2\pi/L$, where L is the length of the cylindrical 3-surface (say a space-time sheet associated with a microtubule). The BE condensates for the resonance frequency photons provide means of communication for the neuron society and could orchestrate microtubules to form a single macroscopic quantum system. One could also consider the possibility that nerve pulse patters are coded to vacuum currents and in turn coded to patterns of coherent photons. In fact, the model of memetic code leads to the identification of nerve pulse/no nerve pulse as Boolean statement true/false. The coding of the nerve pulse patterns to the patterns of vacuum currents of axonal microtubules could occur naturally. The vacuum currents associated with the radial neuronal microtubules could communicate nerve pulse patterns to cell nucleus and the effects of the anesthetics on neuronal microtubules could mean the cutting of this communication line.

A necessary condition for the macroscopic quantum coherence is the phase locking of the vacuum currents associated with different microtubules. Join along boundaries bonds connecting the massless extremals to a larger space-time sheet serving as a common pacemaker could make possible the phase locking.

1.2 Evidence

This picture suggests that microtubules could act as senders and receivers of a coherent light and that visual consciousness should be closely related with the microtubules in accordance with the general philosophy already described. There is indeed some experimental support for identifying the coherent states of photons as associated with vision. It is known that some monocellulars possess elementary vision based on the microtubules [35]. The length distribution of the microtubules in the rods and cones of the eye is concentrated in the region of the visible wavelengths. Insects are known to perceive certain chemical compounds (such as pheromones) by the maser like emission of infrared light by these chemical compounds [43]. Also human nose contains so called vomeronasal organ which seems to give rise to an additional unconscious sense of odors with social and sexual meaning. Interesting hypothesis is that also this vision is based on infrared vision.

There is quite unexpected connection with the phenomenon of sonoluminescence suggesting that liquids contain structures of size of order microtubule diameter and that the highly synchronized light flash emitted in the sonoluminescence results from the condensation of water vapor to liquid involving generation of $k = 149$ space-time sheets from $k = 151$ space-time sheets by p-changing phase transition and subsequent creation of light-like vacuum currents at almost empty $k = 151$ space-time sheets leading to the emission of the flash of coherent light.

An additional support for the quantum antenna hypothesis comes from the quite recently observed anomalous dissociation of water molecules to hydrogen and oxygen in room temperature in presence of catalyst and stirring of the liquid. Usually the reaction is driven by thermal photons at temperature of order 3300 K. A possible explanation is that the Z^0 magnetic flux tubes created by the rotating nuclear Z^0 charge are accompanied by space-time sheets carrying light like vacuum currents generating coherent photons, which in turn drive the dissociation reaction.

Biefeld-Brown effect is one of the oldest poorly understood anomalous effects [56, 57, 58]. What happens is that charged capacitor gains center of mass momentum in the direction orthogonal to the plane of the capacitor plates. Antigravity effect caused by the redistribution of gravitational and/or Z^0 fluxes of the capacitor between various space-time sheets could explain some aspects of the effect. The generation of negative energy space-time sheet with net momentum associated with classical em fields could be also involved with the effect. So called "massless extremals" are optimal candidates for this purpose. This mechanism might be applied by bio-systems to generate coherent motions.

1.3 Quantum antenna hypothesis and brain consciousness

The identification of macroscopic quantum phases possibly serving as quantum correlates of some qualia does not yet help much in understanding brain consciousness. Brain as a neuron society, brain as a music instrument or even entire orchestra and the notion of neural window are metaphors which have served as guidelines in the attempts to guess the general architecture of brain consciousness and might help also the reader to better understand the considerations to follow.

1.3.1 Brain as a neuron society metaphor

The brain as a society of conscious neurons metaphor has surprisingly nontrivial consequences. In particular, a plausible and testable hypothesis for the physical correlates of the sensory qualia becomes possible.

1. Brain as a society of conscious neurons metaphor suggests that our sensory qualia must have a reduction to the neuronal level. For instance, this could mean that our sensory experiences correspond to the sensory experiences associated with the large coherently firing neuron gap junction connected neuron groups in brain.
2. Conscious neurons must be able to communicate their conscious experiences to their fellow neurons. The simplest manner to achieve this is to regenerate the original sensory experience to be communicated by sending a message which creates the stimulus resembling the stimulus giving rise to the original sensory experience.

An attractive idea is that the massless extremals (MEs) associated with the microtubules and other linear structures are for the neuron society what radio receivers and stations are for us. Perhaps the idea about the information society at neuronal level does not look so far fetched if one recalls that a communication based on the genetic code takes place already at DNA-protein level. Furthermore, if Nature has invented a communication by means of a coherent light, it probably has invented also the use of several bandwidths by using several microtubule lengths so that very sophisticated communication systems could exist in brain.

Brain as a society of neurons hypothesis has close relationship to other hypothesis with very similar content. Global workspace hypothesis [21] states essentially that mass media type communication available for large numbers of neurons plays crucial role in the functioning of conscious brain: coherent light is ideal for this purpose. Also brain as hologram idea [25, 30], which is abstracted to neuronal window idea in TGD framework, states that some kind of mass media type communication occurs.

1.3.2 The notion of neural window

The idea of neural window suggests that secondary sensory organs see either the classical em field or the coherent light generated by the mind-like space-time sheets representing the objects of the perceptive field, which can be associated with the primary sensory organ or with the secondary sensory organs in thalamus and cortex. This secondary vision, which could make possible imagination in or all sensory modalities, would be made possible by the coherent photons suffered Bose-Einstein condensation on space-time sheets associated with microtubules or with axons (several space-time sheets might be involved) and serving as wave guides. Massless extremals allow to translate the notion of neural window to the notion of quantum hologram.

1.3.3 Music metaphor

Music metaphor which states that each neuron gives rise to characteristic sensory experience like string of piano gives rise to single note, gives strong constraint on the neuronal window idea. The massless extremal associated with axon corresponds to definite axon dependent frequency. In fact, in the proposed model for the quantum correlates of the sensory qualia [K3] sensory qualia are characterized by some frequency of BE condensed photons besides a pattern of cyclotron frequencies.

A related catching metaphor is to regard groups of parallel axonal microtubules as an orchestra producing light instead of sound with various frequencies. The interior containing the light-like em current would be the instrument and the note produced by single tubule would be a superposition of the frequencies $n\omega_0$, $\omega_0 = 2\pi/L$. The Fourier spectrum of the massless extremal would define the characteristics of the instrument. Of course, in long time scales microtubule could vary its length and achieve more impressive performances than single note samba. A good candidate for the player is the microtubule surface controlling the amplitude of the quantum photon field emitted by the interior by modulating the light-like current in the interior.

1.3.4 Brain as an associative net

The previous metaphors are consistent with the basic view about brain is as associative net such that conscious associations at neural level correspond to conscious experiences of presynaptic neurons associated with the experience of the postsynaptic neuron. The experiences of given neuron is always the same and only its intensity varies so that brain is indeed like a music instrument or orchestra. The intensity of experience is coded by the pattern of nerve pulses. The hypothesis about memetic code states following things. Nerve pulse/no nerve pulse corresponds to true/false Boolean statement; the codons of the memetic code consist of 126 bits and have total duration of order .1 seconds, the duration of our cognitive subself; single bit corresponds to a duration of order one millisecond, the duration of nerve pulse; codons are represented by temporal sequences of cognitive neutrino pairs to which nerve pulse sequences are coded; cognitive neutrino pairs are in turn coded to conscious experiences in many-to-one manner by a unique code analogous to that coding mRNA sequences to polypeptides.

1.4 Relationship of TGD approach with microtubular approach

The role of the microtubules (for a nice introduction see [28]) is believed to be also important for brain consciousness. In TGD framework however microtubules are only one, rather low-lying, although certainly important, level of the self-hierarchy and microtubular consciousness is not expected to correspond to our consciousness directly. In fact, the identification of our subselves (mental images) as 'ELF selves' having as their geometrical correlates topological field quanta of em field with size of Earth, supported by various experimental data about the effects of ELF (extremely low frequency) em fields on brain and correlating our subselves with certain EEG frequencies, could not be philosophically farther from the reductionistic identification of microtubules as seat of our consciousness proposed by Penrose and Hameroff [27].

Fröhlich condensates [34] and microtubular Bose-Einstein condensates of photons have been proposed as the relevant macroscopic quantum phases in the microtubular theory of consciousness. Also in TGD framework macroscopic quantum phases are crucial and serve as quantum correlates of sensory qualia. The basic problem of these theories is how to preserve macroscopic quantum coherence over a time interval of order .1 seconds characterizing our consciousness. In TGD framework the wake-up time of the microtubular selves (time which they are able to stay p-adically unentangled) of about 10^{-16} seconds typically, is not a problem since microtubular selves are not our immediate subselves.

The notion of many-sheeted space-time allows TGD counterparts of both Fröhlich condensates and microtubular photon BE condensates as condensates associated not only with microtubules.

1. Wormhole contacts are unavoidable element of the many-sheeted space-time concept. Wormhole contacts behave in many respects like charged particles and are described by a complex order parameter and it makes sense to speak about wormhole super conductivity. The connection with Fröhlich's condensate comes as follows. Electric fields penetrate from one space-time sheet to another via wormhole contacts carrying quantized fluxes. Thus the normal component of electric field is essentially the density of wormhole charge given by the modulus squared for the order parameter of the wormhole BE condensate. Vacuum polarization of the space-time sheet amounts to the generation of wormhole BE condensates of opposite gauge flux on the two sides of the polarized space-time sheet. In a well defined sense wormhole contact order parameter is square root of the order parameter of Fröhlich condensate.
2. Living matter behaves as liquid crystal and the electret nature of liquid crystals is crucial for many-sheeted ionic flow equilibrium since the weak but coherent electric fields make possible ohmic currents at atomic space-time sheets transforming to supra currents at superconducting space-time sheets.
3. Vacuum gauge fields with non-vanishing gauge currents are a generic phenomenon in TGD and not possible in standard theories. These c-number currents automatically generate quantum coherent states of photons and gravitons via their coupling to the corresponding quantized boson fields. Massless extremals are ideal in this respect since the generation of coherent photons by the light-like vacuum current occurs resonant like manner. Very importantly, massless extremals allow BE condensates of photons in the direction of the light-like vacuum current. This means that massless extremals can serve both as receiving and sending quantum antennae.

1.5 MEs and information molecules

The notion of information molecule is central for the understanding of biological control. There are however several difficult questions related to the notion of information molecule. TGD inspired view about biocontrol and coordination suggests a general answer to these questions and leads to a general model of biological control based on both MEs and information molecules with massless extremals (MEs) serving as actual information carriers initiating self-organization processes whereas information molecules are in the role analogous to that of computer password.

1.6 MEs and quantum holography

One can generalize the original solution ansatz for MEs by introducing what might be called local light cone coordinates for M_+^4 . Boundary conditions for MEs are satisfied if the boundaries of MEs are light-like 3-surfaces, and thus have the same miraculous conformal properties as the boundary of the future light cone. In fact, the light-likeness of the boundaries of M^4 like space-time sheets provide a universal manner to satisfy boundary conditions for field equations.

The superconformal and supercanonical symmetries can be used to generalize the construction of the configuration space geometry to take into account the classical non-determinism of Kähler action. Quantum holography in the sense of the quantum information theory allows to interpret MEs both as receiving and sending quantum antennae as well as dynamical holograms with light-like vacuum currents defining the counterpart of the diffraction grating, and making possible the teleportation of quantum em fields. The superconformal and supercanonical symmetries, which

commute with Poincare symmetries apart from quantum gravitational effects, makes the boundaries of MEs natural seats of supercanonical representations, and since these states are genuine quantum gravitational states defined by statefunctionals in the 'world of worlds', they are expected to be crucial for understanding higher level consciousness.

MEs induce super currents in superconducting magnetic circuits by magnetic induction mechanism, serve as Josephson junctions between magnetic flux tubes, and induce magnetic quantum phase transitions. MEs can generate reference waves or their phase conjugates (time reversals) acting on lower level MEs serving as dynamical holograms. The induced coherent light pattern would act as a control command or its time reversed version. Conjugate reference waves could provide an extremely simple basic mechanism of healing by time reversal allowing the living matter to fight against second law. MEs could read DNA strand to the light-like vacuum current by drifting along it and thus code DNA strand/conjugate strand to a hologram or its phase conjugate in turn acting as a control command or its time reversal. Thus living matter could be regarded as a symbiosis in which MEs control superconducting magnetic flux tubes controlling ordinary matter at atomic space-time sheets via many-sheeted ionic flow equilibrium. DNA would represent the ROM of this system.

1.7 MEs and the notion of conscious hologram

The notion of conscious hologram is the last step in the development of ideas related to bioholograms. The basic challenge is to generalize the notion of the ordinary hologram to that of a *conscious* hologram, about which bio-holograms would be examples. The notion of quantum gravitational hologram is defined at the level of geometric, purely physical existence whereas conscious holograms exist at the level of subjective existence defined by the sequence of quantum jumps and giving rise to the self hierarchy. Of course, these two notions of hologram must be closely related.

The notion of conscious hologram combines the saint and sinner aspects of consciousness to single concept: macrotemporal quantum coherence due to the generation of bound state entanglement and giving rise to co-operation on one hand, and the dissipative self-organization giving rise to Darwinian selection and competition on the other hand.

In nutshell, the notion of conscious hologram follows from the topological field quantization. Classical fields and matter form a Feynmann diagram like structure consisting of lines representing matter (say charged particles) and bosons (say photons). The matter lines are replaced by space-time sheets representing matter (elementary particles, atoms, molecules,...), and virtual bosons are replaced by topological light rays ("mass-less extremals", MEs). Also magnetic flux tubes appear and together with MEs they serve as correlates for bound state quantum entanglement.

The classical fields associated with MEs interfere only at the nodes, where they meet, and one has a hologram like structure with nodes interpreted as the points of a hologram. Thus one avoids the loss of information caused by the interference of all signals everywhere. This aspect is crucial for understanding the role of em fields in living matter and brain. The MEs corresponding to 'real photons' are like laser beams entering the hologram and possibly reflected from it. What is new that the nodes can be connected by 'virtual photon' MEs also analogous to laser beams. Hence also 'self-holograms' with no laser beam from external world are possible (brain without sensory input).

The hologram has a fractal structure: there are space-time sheets at space-time sheets and high frequency MEs propagating effectively as mass-less particles inside low frequency MEs serving as quantum entangling bridges of even astrophysical length. The particle like high frequency MEs induce 'bridges' between magnetic flux tubes and atomic space-time sheets at the receiving end. This makes possible the leakage of supra currents from magnetic flux tubes to atomic space-time sheets analogous to the exposure of film producing hologram. The leakage induces dissipation, self-organization, and primitive metabolism as a cyclic flow of ionic currents between the two

space-time sheets, and thus a Darwinian selection of the self-organization patterns results. Under certain conditions the leakage followed by dropping back to the larger space-time sheet can also give rise to a many-sheeted laser. The low frequency MEs are responsible for the bound state entanglement, macroscopic quantum coherence and co-operation whereas high frequency MEs are responsible for self-organization and competition.

The 3-D vision associated with ordinary holograms generalizes to stereo consciousness resulting in the fusion of mental images associated with the points of conscious hologram [K4].

1.8 Negative energy MEs and bio-control

Negative energy MEs correspond to space-time sheets with a reversed time orientation. These MEs serve as correlates for bound state entanglement. Low frequency negative energy MEs can contain inside them high frequency MEs propagating along them like negative energy particles. The possibility to quantum jump to a higher energy state by generating negative energy ME gives rise to the pay now-let others pay mechanism of metabolism. This quantum credit card mechanism makes the functioning of the living system extremely flexible. The fact that ELF MEs play an important role in living matter forces to consider the possibility of remote metabolism and the transfer of metabolic energy even in the length scale of Earth (7.8 Hz frequency corresponds to Earth's circumference). The small energy dissipation related to 'our' consciousness could perhaps help to solve 'brain's energy crisis' [23] raised by the puzzling observation that the human brain plus body as a whole does not use more energy than smaller brained mammals with a similar body size.

Negative energy MEs are optimal for the realization of intentions. First p-adic ME transformed to a negative energy ME is generated and serves as a geometric correlate of intention. Then quantum jumps of a real system to a higher energy state occurs and in this quantum jumps p-adic ME is transformed to a negative energy ME to take care of the conservation laws.

Right and left brain hemispheres could have different arrows of the geometric time at appropriate p-adic time scales. For instance, negative energy MEs would make possible quantum communications to the direction of the geometric past. The model of non-episodal memory call would involve quantum communication of the question to the geometric past (time-like entanglement and sharing of mental images), and a classical (dissipative) communication of the answer to the geometric future. Negative-positive energy dichotomy could be realized in an extremely wide range of time scales and to explain, besides the basic mechanism of long term memory, also precisely targeted realization of intentions, sensory-motor dichotomy, and biocycles as dissipation-healing cycles.

1.9 MEs and dark matter hierarchy

This chapter has been written much before the emergence of the idea about dark matter hierarchy serving as the source of the long ranged electro-weak and color gauge fields implied by classical. I have left the chapter as such despite the deep insights provided by this vision (for the discussion of this vision see for instance [C6, F6, F9, J1, J2, J6]). Some comments are however in order.

The levels of the dark matter hierarchy are labelled by the value of Planck constant expressible in terms of generalized Beraha numbers $B_q = 4\cos^2(\pi/q)$, $q = m/n$, by the spectrum of complex conformal weights expressible in terms of subsets of zeros of Riemann Zeta, a rational characterizing the effective q-adic topology with which the p-adic topologies corresponding to the primes appearing as factors of the numerator and denominator of the rational number $q = m/n$ are consistent, and an algebraic extension of p-adic number field in question. It is clear that the visible matter is in this framework only a tip of an iceberg. A good metaphor for TGD Universe is as an inverted Mandelbrot fractal for which the introduction of longer length scale resolution

meaning bird's eye of view makes visible new large scale structures in every new zoom without limit.

An attractive hypothesis is that particle is characterized either by the integer m or n (nothing to do with q above) depending on whether it corresponds to a positive or negative energy particle, that two positive (negative) energy particles can have direct interactions if the integers m (n) associated with them have common prime factors, and that these primes characterize the elementary particles exchanged in the interactions.

As a rule of thumb one can say that the values of \hbar coming as $\lambda^k \hbar$, $\lambda = nv_0$ $v_0 \simeq 2^{11}$ define dark matter hierarchies such that higher level particles decay to lower level particles by de-coherence phase transition reducing the value of \hbar by $1/\lambda$. This hierarchy means for a given particle a hierarchy of zoomed up Compton lengths and times making possible macroscopic quantum coherence by the overlap criterion of space-time sheets having sizes of order Compton length.

The criterion for the phase transition to larger \hbar phase at the lowest level is that the interaction strength αQ^2 for particles of charge Q and gauge coupling strength α satisfies $\alpha Q^2 \geq 1$ and implies the increase of \hbar by $\hbar \rightarrow Q^2 \alpha \hbar / v_0$ implying the reduction of the interaction strength as $Q^2 \alpha \rightarrow v_0$.

MEs can be regarded as space-time correlates for a hierarchy of gauge bosons characterized by different values of Planck constant and the de-coherence phase transition would naturally correspond to the decay of MEs to smaller space-time sheets. The ordinary laser light cannot be regarded as a large \hbar phase, which de-coheres to ordinary photons before the interaction with ordinary matter. Very general consistency arguments lead to the conclusion that dark matter and dark MEs must correspond to λ^k -fold ($k > 0$) coverings of M^4 locally ($\hbar(k) = \lambda^k \hbar_0$, $\lambda \simeq 2^{11}$) whereas ordinary laser light would correspond to $k = 0$.

2 Massless extremals

The so called massless extremals are very general solution of field equations associated with the minimization of Kähler action parameterized by several arbitrary functions. The purely TGD:eish feature of the massless extremals is the presence of light-like currents generating coherent states of photons and gravitons. These features suggest that massless extremals might have important role in bio-systems.

2.1 Massless extremals as general solutions of field equations

Let $k = (k^0, k^3, 0, 0)$ be a light like vector of M^4 and $u = u(m^1, m^2)$ arbitrary function of the Minkowski coordinates m^1 and m^2 in the plane orthogonal to the direction of the 3-vector $(k^3, 0, 0)$ associated with k . The surfaces defined by the map

$$s^k = f^k(k \cdot m, u) , \quad (1)$$

where f^k and u are arbitrary functions define massless extremals. They describe the propagation of massless fields in the direction of k : the fields are periodic with a period $\lambda = 2\pi/k$ so that only k and its integer multiples are possible wavevectors. The polarization associated with various induced gauge fields depends on the position in (m^1, m^2) -plane and is in the direction of the gradient of u . Field equations involve tensor contractions of the energy momentum tensor and gauge current but these are proportional to kk and k respectively and vanish by the light-likeness of k . Linear superposition holds true only in a restricted sense since both the propagation direction and the polarization direction in each $(m^1, m^2) = const$ plane is fixed.

What is remarkable that these solutions are not solutions of the ordinary Maxwell equations in vacuum: Kähler current density J_K is in general non-vanishing(!) and proportional to the light

like four-momentum k . As a consequence, also a light-like electromagnetic current is in general (but not necessarily) present. The interpretation of the em current J as charged elementary particle current is impossible and the correct interpretation as a vacuum current associated with the induced gauge fields. The finite length of the microtubule plus the requirement that the total vacuum charge vanishes, implies that the Fourier decompositions of the massless fields contain only integer multiples of the basic four-momentum k . The direct detection of the light-like vacuum current inside a microtubule would provide strong support for TGD.

The physical importance of these extremals is suggested by the fact they are in certain sense elementary particle like objects: in fact, the original interpretation was as a model for the exterior space-time of a topologically condensed massless particle. The solution set is also very general involving several arbitrary functions. Although the minimization of the Kähler action favors the formation of Kähler electric fields, massless extremals might well appear as space-time sheets of the effective space-time. These space-time sheets should not contain ordinary charges since their presence implies a transition to the Maxwell phase described in an excellent approximation by the ordinary Maxwell electrodynamics.

Rather remarkably, massless extremals are also solutions of the field equations associated with the low energy effective action. This holds true in the absence of the topologically condensed matter, phenomenologically described using external currents. For instance, the term

$$(T_{\#}^{\alpha\beta} - \frac{1}{16\pi G} G^{\alpha\beta}) D_{\beta} \partial_{\alpha} h^k ,$$

where $\#$ refers to the topologically condensed matter, reduces to

$$\frac{1}{16\pi G} G^{\alpha\beta} D_{\beta} \partial_{\alpha} h^k ,$$

and vanishes identically because Einstein tensor is light like so that contraction with the second fundamental form vanishes. The vanishing of these terms in presence of matter is not possible since the gauge currents and energy momentum tensor associated with the topologically condensed matter are not light-like. Thus massless extremals correspond to vacuum space-time sheets with respect to the ordinary matter. Massless extremals can however interact with the ordinary matter via $\#$ contacts.

The fact that vacuum em current and vacuum Einstein tensor do not in general vanish, implies that massless extremals serve as sources of coherent photons and gravitons. It is not very economical to maintain BE condensates all the time. In 'dormant' states microtubules could correspond to ME:s with vanishing em fields but non-vanishing Z^0 fields or even vacuum extremals of the effective action with one-dimensional CP_2 projection and having vanishing classical gauge fields. Massless extremals can also reduce to vacuum extremals of the Kähler action in the case that the CP_2 projection is, in general two-dimensional, Legendre manifold of CP_2 . Also in this case massless extremals are however non-vacuum extremals of the effective action.

2.2 About the electro-weak and color fields associated with massless extremals

Space-time sheets carrying em fields carry usually also Z^0 and W fields and it is not possible to speak about em or Z^0 type MEs. It is however possible to speak about neutral and W MEs. The CP_2 projection of ME is 2-dimensional and in a special case it reduces to a geodesic sphere. There are two kinds of geodesic spheres in CP_2 .

1. For space-time sheets for which CP_2 projection is $r = \infty$ homologically non-trivial geodesic sphere of CP_2 one has

$$\gamma = \left(\frac{3}{4} - \frac{\sin^2(\theta_W)}{2} \right) Z^0 \simeq \frac{5Z^0}{8} .$$

The induced W fields vanish in this case and they vanish also for all geodesic sphere obtained by $SU(3)$ rotation.

2. For homologically trivial geodesic sphere a standard representative is obtained by using for the phase angles of standard complex CP_2 coordinates constant values. In this case induced em, Z^0 , and Kähler fields vanish but induced W fields are non-vanishing. This holds also for surfaces obtained by color rotation. Hence one can say that for non-vacuum extremals with 2-D CP_2 projection color rotations and weak symmetries commute.

The MEs corresponding to these two geodesic spheres could be called neutral and W MEs and they carry color fields for which the color group $SU(3)$ reduces to some of its $U(1)$ subgroups. Quite generally, the holonomy algebra of color group is Abelian since the induced color field is of the form $g_{\alpha\beta}^A \propto H^A J_{\alpha\beta}$, where H^A denotes color Hamiltonian.

Neutral MEs are excellent candidates for mediating EEG type communications from the biological body to the magnetic body whereas charge entanglement induced by W MEs would be ideal for the realization of motor actions of the magnetic body by generating first superposition of exotically ionized states of atomic nuclei entangling magnetic and biological body [M3]. State function reduction would lead to an exotically ionized state accompanied by dark plasma oscillation pattern. By Faraday law this pattern would induce electric fields at the space-time sheets containing ordinary matter which in turn would generate ohmic currents leading to various physiological effects.

MEs are excellent candidates for the space-time correlates of laser beams. Dark matter hierarchy implies that also MEs can be classified by the level of the dark matter hierarchy involved. A very general argument leads to the conclusion that dark space-time sheets, in particular MEs, at the k^{th} level of the dark matter hierarchy correspond to space-time sheets defining λ^k -fold coverings of M^4 (recall that one has $\hbar(k) = \lambda^k \hbar_0$ and $\lambda \simeq 2^{11}$) [C6, M3]. $k = 0$ MEs would correspond to the ordinary laser light.

2.3 How massless extremals generate coherent states of photons?

ME:s can be in 'dormant' or active state according to whether the em current associated with the ME is vanishing or not. In active state ME:s generate Bose Einstein condensate type state for ordinary photons. This means in TGD context the emission of (topological) vapor phase photons (CP_2 type extremals), which can condense on other condensate levels. ME:s generate gravitonic BE condensate and the possible biological role of this condensate will be discussed later.

Assuming that the coupling of quantized photon field to the massless extremal is given by regarding the massless extremal as a classical background field one obtains QED with a light like source J^α :

$$\begin{aligned} D_\beta F^{\alpha\beta} &= eJ^\alpha , \\ J^\alpha &= Jk^\alpha . \end{aligned} \tag{2}$$

The system is equivalent with an infinite number of harmonic oscillators each driven by a harmonic external force and a basic exercise in the quantum mechanics shows that the solutions of the field equations give the new oscillator operators as sums of free oscillator operators plus c-number term, which is essentially the Fourier component of the light like current in the direction of the polarization.

In the limit that ME has infinite duration and is a cylindrical structure of finite length L (that is microtubule) one has for $J \propto \sin(k_z(t-z))$

$$\begin{aligned}
a^\dagger(p) &\rightarrow a^\dagger(p) + g(p) \ , \\
g(p) &= \sum_n \delta(p^0, k_n^0) K(p, k_n) J(k_n^z, p_T) \ , \\
K(p, k) &= \epsilon(p) \cdot k \frac{1}{i(p_z - k_z)} (\exp(ip_z L) - 1) \ , \\
k_n &= nk_0 = \frac{n2\pi}{L} (1, 1, 0, 0) \ .
\end{aligned} \tag{3}$$

Here p denotes the momentum of the photon and k the 4-momentum associated with the Fourier component of a light-like current. $\epsilon(p)$ denotes the polarization of the photon. $J(k_n^z, p_T)$ is essentially the 3-dimensional Fourier transform of the scalar function J . The infrared behavior of $J(k_z, p_T)$ as a function of the transversal momentum p_T can be deduced from the fact that the transverse dimension of the microtubule is small (about 25 nm) as compared to $1/p_T$ so that the Fourier component is in good approximation independent of p_T .

For the frequencies present in the Fourier decomposition of the massless extremal, the ordinary oscillator vacuum is transformed to a coherent state in the corresponding Fourier mode of the quantized photon field. The essential point is that the wave vectors of the radiation field and massless extremal are nonorthogonal. The radiation pattern resembles the ordinary antenna pattern associated with an oscillating current $J(t) = \exp(i\omega t)$ in that the intensity of radiation vanishes at angles $\theta = \pi/2$ and $\theta = 0$. For $J \propto \sin(k_z(z-t))$ $|K|^2$ has maxima for $\theta = 48.6$ degrees and 131.4 degrees. For an ordinary dipole with $J = \sin(\omega t)$, $\omega = 2\pi/L$ the radiation pattern is concentrated at angles $\theta \geq 40$ degrees with maximum and 69.3 degrees and 110.7 degrees.

A more complicated situation corresponds to a group of several massless extremals (say microtubules). If massless extremals are parallel and have same length the previous expression generalizes with superposition of terms

$$g(p) \rightarrow \sum_n \exp(i\phi_n) \exp(ip_z z_n) \exp(ip_T \cdot x_T) g_n(p) \ . \tag{4}$$

The phase ϕ_n is the phase difference between n :th light like current with respect to some reference current. If the positions of microtubules and/or phases of the individual light like currents are suitably chosen then various terms interfere constructively and macroscopic quantum coherence is obtained at resonant frequencies. Suffice it so say that the needed timing is extremely accurate: less than 10^{-12} seconds! Since p_z is small rather larger transversal distances are allowed by the requirement of constructive interference. In a more general situation also the orientations of microtubules can vary in certain limits. Note that light-like energy momentum generates also gravitonic BE condensates at preferred frequencies.

2.4 Massless extremal is accompanied by a Bose-Einstein condensate of parallel photons

The interaction Lagrangian describing the interaction of photon field with the light-like vacuum current does not couple to the photons collinear with the vacuum current (light-like wave vector has vanishing length squared). Therefore the ground states of the system are degenerate since one can add to any coherent state generated by the vacuum current any number of photons collinear

with the vacuum current and topologically condensed inside the massless extremal. This means Bose-Einstein condensation in collinear degrees of freedom.

Collinear Bose-Einstein condensates of photons are crucial for the model of the quantum correlates of the sensory qualia. Sensory quale is characterized partially by the BE condensate of photons associated with the massless extremal parallel to the axon. The existence of the BE condensate makes possible induced emission. For instance, Josephson currents generate photons with frequencies which are multiples of the Josephson frequency. If the potential difference in Josephson junction equals to a multiple of the cyclotron frequency of some superconducting ion, the current flows resonantly in the sense that Josephson current serves as a harmonic perturbation generating quantum jumps and gives rise to a large dissipative current and also quantum jumps in either superconductor. Since the emission rate for photons by the current is proportional to N^2 , where N is the number of photons already in the state, the presence of the BE condensate of photons with this frequency amplifies the emission rate. This kind of resonance mechanism is assumed in the model of sensory experience since it elegantly explains why given neuron corresponds to single quale. Since the potential difference over the Josephson junction can correspond to only single cyclotron frequency, the dominance of single quale is unavoidable even when all macroscopic quantum phases are present.

The existing BE condensate increases the probability of topological condensation of coherent photons generated by other massless extremals to the massless extremal. This mechanism could provide inter-neuronal communication mechanism and realize the metaphor about brain as a society of neurons, the notion of neuronal window idea and also give a more precise content to the music metaphor. In particular, neurons far away from each other could communicate using wavelengths in a narrow wave length range by this mechanism.

The wave vectors of the photons are multiples of $k = \pi/L$. This means that the length of the massless extremal correlates with the maximal allowed wavelength. For ELF photons associated with EEG frequencies of order 10 Hz the length of massless extremal is of order Earth's circumference. This suggests that more general massless extremals with a topology of torus instead of linear topology could characterize the topological field quanta of ELF fields. It is however impossible to say, whether the field equations allow more general solutions resembling massless extremals.

2.5 MEs and exotic representations of p-adic Super Virasoro

The exotic representations of p-adic Super Canonical and Super Virasoro algebras forming excellent candidate for a hierarchy of lifeforms are associated with MEs [M2, K3]. For these representations mass squared of the particle is proportional to Virasoro generator L_0 having integer valued mass spectrum. When the value of the p-adic mass squared is power of p : $M^2 \propto p^n$, $n = 1, 2, \dots$, the real counterpart of the mass squared is extremely small since it is proportional to $1/p^n$ in this case. These states are generated by a subalgebra of Super Canonical Algebra in case of MEs since the conformal weight of the vacuum state vanishes (for ordinary elementary particles conformal weight of the vacuum is negative integer). Thus very special representations of p-adic Super Virasoro algebra are in question.

The degeneracy of states (number of states with same mass very small squared) is enormous for the physically interesting values of p-adic prime p . This means that these states provide huge negentropy resources. Thus exotic Super Virasoro representations be interpreted as quantum level articulation for the statement that TGD Universe is quantum critical quantum spin glass. Exotic Super Virasoro representations clearly provide an excellent candidate for an infinite hierarchy of life forms. These lifeforms are labelled by three integers (k,m,n): physically interesting primes correspond to $p \simeq 2^{k^m}$, whereas k prime and m are integers, and the power n appearing in $M^2 \propto p^n$. It is these lifeforms which make mind-like space-time sheets living creatures and these lifeforms emerge already in elementary particle length scales and become increasingly complex when the p-

adic length scale increases. Life can be regarded as a symbiosis of these lifeforms. These lifeforms ('mind') interact with each other and ordinary matter via the classical gauge fields associated with MEs. A natural hypothesis is that the quantum phase transitions of the macroscopic quantum phases for the particles of the exotic Super Virasoro representations formed in classical fields of MEs (mind-like space-time sheets) give rise to some (but not all) qualia.

3 Microtubules as quantum antennae

3.1 Linear structures as quantum antennae

The many-sheeted space-time concept of TGD indeed allows almost vacuum space-time sheets and these space-time sheets might be crucial for the understanding of the bio-systems. For instance, the weak interaction of these space-time sheets with the ordinary space-time sheets containing matter could provide representations of the external world in the physical properties of the almost vacuum space-time sheets. In particular, mind-like space-time sheets having finite time duration could generate coherent light and coherent light might make communication possible between mind-like space-time sheets and realize the idea of neuronal window discussed briefly in introduction [H7]. The mind-like space-time sheets associated with various linear structures are especially natural candidates for massless extremals serving as quantum antennae. Bio-systems are full of linear structures and the mind-like space-time sheets associated with microtubules, DNA and protein molecules are the most obvious candidates for quantum antennae.

3.2 Are microtubules accompanied by massless extremals?

The interior of the microtubule is by its cylindrical symmetry an ideal place for ME:s whereas the axonal cell membranes must correspond to 'Maxwell phase', where ordinary Maxwell equations are satisfied in a good approximation but also a separate space-time sheet is possible candidate for a massless extremal. The function $u(x, y)$ appearing in the general solution is most naturally $u(x, y) = \sqrt{x^2 + y^2}$ for microtubules implying radial polarization. The explanation of the macroscopic quantum coherence in the brain would provide crucial support for the TGD based world picture since massless sources and vacuum currents are not possible in the ordinary QED.

There is analogy with the super radiance phenomenon [49]: in this case however the photons radiated by the microtubule waveguide have momenta parallel to the microtubule so that the mechanism leading to the formation of the macroscopic BE condensate remains to be understood. The theories associating coherent photons with the microtubules typically assume that the coherent photons reside inside the microtubules: this leads to problems since Uncertainty Principle; the direct study of Maxwell equations also suggest that photons should have very large transversal momenta corresponding to the transversal dimension of the microtubule of order 10^{-8} meters. In TGD this difficulty is avoided since only the *sources* of the coherent photons are restricted into the interior of microtubules whereas the photons can exist in vapor phase and condense on various space-time sheets of the topological condensate.

The necessary condition for the formation of the coherent states in the presence of the matter is that ME condensate level does not contain ordinary gauge charges. If ME corresponds to a larger space-time sheet 'below' the space-time sheet containing the ordered water, this requires that the gauge charges of the condensed matter do not flow to the interior of the ME space-time sheet. This is achieved if the condensed particles combine by join along boundaries bonds together to form a net like structure so that gauge fluxes can run along the join along boundaries bonds to the boundary of the ME region, where they can flow down to the ME condensate level. Ordered water is a good candidate for this join along boundaries condensate. Microtubules are known to be surrounded by ordered water and also the interior contains ordered water. The axial electric

polarization of the microtubular surface suggests that there is a longitudinal electric gauge flux at the condensate level of the ordered water running to/from the ME condensate level at the ends of the microtubule. The wave lengths appearing in the Fourier expansion of J are of form L/n , L being microtubule length.

In the active mode microtubule acts as a quantum antenna creating quantum coherent light unlike the ordinary antenna, which creates incoherent light. Also waveguide mode is possible for the topologically condensed photons but antenna mode is crucial for the generation of the macroscopic coherent states. The following argument suggests that the dielectric properties of the microtubule surface can change the antenna pattern somewhat. The dipoles of the microtubule are known to be parallel to the axis of the microtubule. The interaction energy of a dipole p with the radiation field is proportional to the quantity $p \cdot E = pE \cos(\theta)$ so that the effect of the dielectric is largest, when the wave vector of the photon is orthogonal to the axis of the microtubule. As a consequence, the dipole pattern should concentrate in the forward direction.

The em current in the interior transforms ordinary QED vacuum to a coherent state in the resonating Fourier modes of the photon field. In ME mode the resonance energies come as multiples of $E_0 = 2\pi/L$ and wavelength L/n , where L is the length of the microtubule whereas in VE mode the spectrum is continuous. Biophotons [52] with energy of order one eV might be regarded as evidence for BE states associated with the shortest microtubules (such as centriole and basal bodies). The average length of the neuronal microtubules is about $10^{-5} m$ and corresponding IR radiation is more energetic than thermal IR radiation with a wavelength of order $10^{-4} m$.

In the ME mode the resonances at energies $E_n = nE_0$, $E_0 = 2\pi/L$, provide ideal communication channels. Microtubules with different lengths provide independent communication channels so that very effective communication in principle becomes possible. This process could orchestrate axonal microtubules as well as the microtubules belonging to different neurons to form a larger macroscopic quantum state. An optimal performance is obtained if the microtubules belonging to same group are parallel and their lengths are quantized with a common multiple. The microtubules of the neighboring neurons indeed tend to be parallel. Axonal microtubules are also parallel whereas the microtubules inside the ordinary cells are in radial configurations. The grey matter in brain has a columnar structure so that axonal microtubules tend to be in the direction of the columns: this should favor the formation of a quantum resonance between different microtubules. Furthermore, the model of [50], described in Tuscon II, for the microtubule interactions predicts that the microtubules of even far away neurons are parallel. The average length of the neuronal microtubules is about $10^{-5} m$ and it is known that the response of 3T3 cells to weak IR radiation is maximum at this wavelength. Neurons could be able to tune their microtubules to the desired infrared stations by controlling their orientation and length. The upper bound about 10^{-4} meters for the length of the axonal microtubules can be understood: for the longer microtubules the thermal IR radiation becomes important and makes communication impossible. In long axons this problem is avoided by joining shorter microtubules in series via Microtubule Associated Proteins (MAPs).

Since the time scale for the change of the tubulin polarization is of order 10^{-10} seconds and the period of the IR radiation is of order 10^{-13} seconds, amplitude modulated IR transmissions are possible. The mechanism of the amplitude modulation could be simply a change of the microtubule interior gauge field from active to dormant ME mode. Amplitude becomes vanishing if this field becomes ordinary sourcefree em field or Z^0 field. IR transmissions could be based on some kind of binary code resembling genetic code. There is indeed concrete proposal of Koruga for this code motivated by the geometric structure of the microtubule surface. [37, 28]. One possibility is that the propagating modes of dipole and conformational oscillations perform elementary AM modulations. These modes could correspond to elementary language expressions at the level of the microtubules.

Microtubules can also absorb photons coming from an external source at resonance energies.

If Bose-Einstein condensate of N photons in some mode is present, the absorption probability is amplified roughly by the factor N^2 as shown in appendix. This suggests that microtubules containing BE condensate of photons in some mode are able to 'see' in some elementary sense. Of course, receiving antenna containing the BE condensate need not be microtubule. Centrioles (T shaped pair of microtubules inside animal cells) could provide cell with infrared eye and there is experimental evidence for this in the case of monocellular organisms [35]. Also the radial microtubules could have elementary sense of vision. Note that all eucaryotic cells have radial structure of microtubules in their cytoskeleton.

3.3 How macroscopic quantum coherence is generated?

The big problem is the creation of constructive interference between the coherent states associated with different microtubules. The problem looks exceedingly difficult: microtubules should be able to tune up the frequency and phase associated with light like current inside microtubule with those of other microtubules contributing to the coherent state. Frequency tuning, or equivalently length tuning, involves time scales smaller than 10^{-13} seconds in case of infrared light associated with longest microtubules. The simplest solution to come into mind is that there exist some pacemaker keeping the microtubules in rhythm. One can imagine several mechanisms important for the tuning, each involving the special properties of the TGD:ish space-time crucially.

3.3.1 Topological field quantization

TGD:ish space-time surface decomposes into regions characterized by vacuum quantum numbers, which are frequencies and integers related to the time and angle dependences of the phase angles associated with the two complex CP_2 coordinates. Typically one has $\Phi = \omega t + \text{Fourier expansion}$ so that space-time surface vibrates with frequency ω . This vibration is an ideal candidate for a pacemaker for the physical systems inside a given space-time region. In fact, the vacuum quantum numbers characterize partially also the order parameter of a super conductor. Vacuum frequencies could also be special frequencies for the Maxwell fields.

The increased understanding about topological field quanta as classical and quantum coherence regions of em field is consistent with and generalizes this view. When topological field quanta are joined by join along boundaries bond generated in quantum jump they fuse to form a larger region of classical and quantum coherence. This suggests a general mechanism for how various axons/microtubules can generate phase coherent em fields. What is needed is that there is larger space-time sheet connected by join along boundaries bonds to the massless extremals associated with various axons/microtubules. This larger space-time sheet is most naturally the geometrical counterpart of higher level self so that consciousness is what creates synchrony rather than vice versa!

A further important aspect in the generation of synchrony is self-organization. The subsystems of self quantum self-organize and end up to asymptotic self-organization patterns selected by dissipation. These patterns are simple and typically involve spatially repeating patterns and synchronous oscillations (Benard flow is simple example of this). It is consciousness which implies synchrony and coherence whereas in standard approaches to quantum consciousness synchrony and coherence are believed to be prerequisites for consciousness.

3.3.2 Phase locking for the system of Josephson junctions

Japanese physicist Yoshiki Kuramoto from the University of Kyoto has shown that the solutions of the differential equations describing Josephson junctions tend to a state in which there is single collective oscillation frequency. A.T. Winfree from the University of Arizona has shown that a

phase transition to single collective oscillation frequency analogous to the freezing of liquid occurs in this kind of system.

The solutions of the differential equations describing Josephson junctions model quantum self-organization and synchronization can be interpreted as an instance of the selection of asymptotic self-organization patterns selected by dissipation and occurring always in quantum self-organization. Of course, the quantum jumps can occur only provided the system of Josephson junction belongs to a system having self so that consciousness is again prerequisite for synchrony rather than vice versa! In any case, the fact that entire brain is hierarchy of space-time sheets such that the space-time sheets at various levels are connected by Josephson junctions makes this result rather encouraging.

3.3.3 Gap junctions and MAPs

As noticed, the formation of the join along bonds is the basic prerequisite for the formation of the macroscopic quantum systems. The so called gap junctions can be regarded as join along boundaries bonds between the cell membranes (understood as 3-surfaces in TGD picture) of the neighboring cells. They could have a key role in synchronous firing of neuron groups. Gap junctions could also force the vacuum quantum numbers of the neighboring cell membranes to be identical as well as provide the bridges for the propagation of the Maxwell type fields between neighboring cells. It is known that the coherently firing neuron groups in brain possibly responsible for the generation of sensory experience are gap junction connected. It is not however obvious whether gap junctions have anything to do with the synchronizing of the vacuum currents: the difference between the time scales involved is indeed huge. Also Microtubule Associated Proteins could act as join along boundaries bonds guaranteeing quantum coherence between microtubules inside same cell.

3.4 Are nerve pulse patterns coded into vacuum currents and coherent light?

It has turned out that TGD based model for memetic code leads to the same interpretation of nerve pulse patterns as suggested by neuroscience. Nerve pulse/no nerve pulse corresponds to true/false Boolean statement or 1/0 value of binary digit. Fundamental coding of nerve pulse patterns is the coding into temporal sequences of cognitive neutrino pairs associated with cell membrane such that the spin of the cognitive antineutrino codes for true/false (1/0). Of course, bits could code also for binary digits in the binary expansion and code for the intensity of the primitive sensory experience associated with the neuron. It is natural to ask whether nerve pulse patterns could be also coded to some other representations. Light-like currents are indeed optimal in this respect.

The dependence of the light-like current on the longitudinal coordinate of massless extremal is arbitrary and therefore light-like current provides ideal tool of classical communication of information with light velocity as well as coding of this information to coherent light received by other massless extremals. The first bio-application to come into mind is that the instantaneous nerve pulse patterns propagating along the axon could be coded into the pattern of the vacuum current. The velocity of propagation for the nerve pulse pattern is extremely small as compared to light velocity but this is not a problem if the coding takes place in the region where nerve pulses are generated. What happens is that same temporal pattern of pulses propagates with different velocities. This coding in turn implies coding of nerve pulses to coherent states of photons and in principle the communication of nerve pulse pattern to other neurons.

The relationship between memetic and genetic code is that between two hierarchy levels of a computer program. This suggests that nerve pulse patterns representing memetic codons could serve as transcription factors at gene level. This requires the communication of nerve pulse patterns to nucleus. Even more, communication mechanism must treat different presynaptic inputs as

different inputs. Modulation of the vacuum current could make possible communication of the nerve pulse patterns to the cell nucleus along the massless extremals associated with the radial microtubules which in case of neurons have direct contact with the cell membrane. The fact, that some anesthetics seem to affect microtubules and that some brain diseases involve changes of microtubules could also be explained as a breaking of the cell membrane-nucleus communication link. That this kind of communication link might exist is suggested by the fact that ELF em fields have direct effect on genetic expression [29].

4 Masless extremals and information molecules

The notion of information molecule is central for the understanding of biological control. There are however several rarely asked questions related to the notion of information molecule: in particular, the phenomenon of pleiotropy is not easy to understand on basis of pure chemistry [24]. TGD inspired view about biocontrol and coordination suggests a general answer to these questions and leads to a general model of biological control based on both MEs and information molecules with massless extremals (MEs) serving as actual information carriers and information molecules having a role analogous to that of computer passwords.

4.1 Questions about information molecules

Central nervous system (CNS), endocrine system and immune system are three basic systems involved with bio-control and -communication. The work of Candace Pert and other neuroscientists has led to a general notion of information molecule described in popular manner by Candace Pert [26]. Neural transmitters and modulators associated with CNS are only special cases of information molecules. Also neuropeptides and various hormones are involved. It has become clear that emotions are closely related with the activity of information molecules and that both brain, endocrine system and immune system communicate intensively with each other. One could regard even brain as a big gland. Of course, one could also consider various glands and organs as mini-brains.

The interactions of the information molecules involve the formation of receptor-information molecule complex either at cell surface or in the cell plasma inside cell. Receptor-information molecule complex inside cell can move to genome and induce gene transcription. In case that the complex is formed at the surface of cell, second messenger action is involved. One can also speak about N:th messenger action. There are many poorly understood aspects related to the mechanisms of information molecule action [24].

1. There are only few second messenger pathways and relatively few receptors but large number of different functions. This phenomenon is known as pleiotropy or multi-functionality. For instance, given second messenger causes different effects depending on the hormone that activated it (the phenomenon is somewhat analogous to the phenomenon in which message can be understood in several manners depending on the state of receiver). At purely chemical level the problem is how second messenger knows what hormone activated it? In steroid action the complex formed by information molecule and receptor in turn activates some gene. Now the question is: How the activated RNA polymerase knows which gene has to be activated? Pleiotropy appears also at level of hormones. Same hormone can have multiple effects and the border between hormone, neuropeptide or even neurotransmitter is unclear. For instance, hormone which by definition transmits long distance communications, can have effects in nearby cells and thus acts like a neuropeptide. How hormone knows what function it must perform? Also drugs and treatments can have different effects and side effects.

2. There is also functional redundancy: the same function is performed by several second messenger molecules. For instance, glucagon, growth hormone, adrenaline and corticosteroids elevate glucose levels. This suggests that there is deeper level of communication involved and that second messenger molecules are more like computer passwords than subprogram calls. Now the question is: What these subprogram calls do correspond physically?
3. Biological functions can be initiated also in nonchemical manner. The phenomena of healing by touch and the effects of meditation and biofeedback are examples of biological self-organization processes are initiated in nonchemical manner. Even other treatments like massage, acupuncture or meditation can decrease or inhibit pain. These observations suggest that chemical level is not the deepest level involved with biological functions and the question is: What is this deeper control level?

Simple lock and key mechanism cannot provide answer to the questions raised above. Rather, computer password might provide better metaphor for the second messenger action whereas receptor-information molecule complex would effectively generate subprogram call perhaps carried by the second messenger molecule or possibly broadcasted. It seems that information molecules act more like signs or symbols rather than being purely chemical agents. These symbols are interpreted by cell level intelligences and the interpretation depends on context.

4.2 A model of biological self-organization based on quantum antenna hypothesis

The view that self hierarchy is present already at molecular level and realized in terms of MEs provides rather straightforward interpretation of pleiotropy and redundancy. The phenomenon of pleiotropy suggests there is nonchemical communication between receptor-peptide complex and cell nucleus. The most natural TGD inspired candidate for the communication is wake-up of genome subself by general wake-up mechanism in which classical em field associated with ME induces quantum jumps leading to quantum phase transition which could correspond to the transcription process. The almost-determinism of the transcription process would be due to the Darwinian selection caused by dissipative effects.

These considerations suggest that information molecule-receptor complex could generate ME carrying classical gauge fields and vacuum current. Vacuum current is excellent candidate for coding the information and can lead to a generation of coherent light.

1. The first possibility is broadcasting. The ME associated with information molecule receptor-complex acts as active quantum antenna and activated structure, say genome, serves as a passive quantum antenna receiving the coherent light. Classical fields and/or coherent light would induce quantum jumps serving as seeds of quantum phase transitions leading to a wake-up of conscious and self-organizing subself inside receiver.
2. Alternatively, second messenger molecule could carry the information carrying ME with itself as a genuine message inducing the self-organization process in, say, genome.

A natural hypothesis is that the states of the exotic Super Virasoro representations define the macroscopic quantum phases in question: the reason is that these representations are present in all length scales. The information molecule-receptor pair corresponds to a definite frequency, or more generally, combination of frequencies, coding the corresponding function. For instance, genes might be coded to harmonics of Super Virasoro frequencies associated with various p-adic length scales. All information molecule-receptor combinations initiate some function determined by these frequencies and pleiotropy emerges as a basic prediction of the model. Second messenger pathway is like a password to computer, universal key, together with the frequency or even entire ME

specifying the function in question: this initiates the desired self-organization process waking-up proper subself.

These ideas suggest the following general framework for understanding biological self-organization.

1. Biological programs consist of self-organization patterns generated by classical gauge fields associated with MEs at specific resonance frequencies inducing quantum jumps leading to quantum phase transitions. These resonance frequencies serve as names of the genetic sub-programs. Messenger molecules in turn serve in the role of computer passwords.
2. Self-organization processes are associated with MEs and generated by special frequencies, which could be harmonics of the fundamental frequencies associated with various exotic Super Virasoro representations. For instance, combinations of harmonics of various Super Virasoro transition frequencies could define 'name of gene'. The fact that these frequencies are constants of Nature means that the model is immediately testable. Of course, also other transition frequencies, in particular, magnetic and Z^0 magnetic transition frequencies, could be important.
3. The ability of a biological system to act effectively like a deterministic computer is due to the Darwinian selection of the asymptotic patterns of self-organization caused by dissipation in systems which are feeded by energy.
4. The four-dimensionality of this self-organization process is also important element. The frequency of ME defines time scale $T = 1/f$ which defines the duration of biological chronon. With this interval of geometric time entire 4-dimensional space-time surface changes in self-organization process.

5 Evidence for quantum antenna hypothesis

In the following some evidence for quantum MEs and quantum antenna hypothesis is discussed. It must be emphasized that there is also other evidence discussed in other chapters of the book (for instance, see the chapters [K3, M2, M4, M5].

5.1 TGD inspired model for sonoluminescence

Sonoluminescence [42] is a peculiar phenomenon, which might provide an application for the hydrodynamical hierarchy. The radiation pressure of a resonant sound field in a liquid can trap a small gas bubble at a velocity node. At a sufficiently high sound intensity the pulsations of the bubble are large enough to prevent its contents from dissolving in the surrounding liquid. For an air bubble in water, a still further increase in intensity causes the phenomenon of sonoluminescence above certain threshold for the sound intensity. What happens is that the minimum and maximum radii of the bubble decrease at the threshold and picosecond flash of broad band light extending well into ultraviolet is emitted. Rather remarkably, the emitted frequencies are emitted simultaneously during very short time shorter than 50 picoseconds, which suggests that the mechanism involves formation of coherent states of photons. The transition is very sensitive to external parameters such as temperature and sound field amplitude. In the following only the rough hydrodynamical characteristics of the phenomenon are considered from the point of view of p-adic length scale hypothesis. Also an attempt to understand the mechanism behind quantum coherence in terms of light-like vacuum currents associated with massless extremals is made.

5.1.1 Sonoluminescence and hydrodynamic hierarchy

A plausible explanation for the sonoluminescence is in terms of the heating caused by shock waves launched from the boundary of the adiabatically contracting bubble [42]. The temperature jump across a strong shock is proportional to the square of Mach number and increases with decreasing bubble radius. After the reflection from the minimum radius $R_s(min)$ the outgoing shock moves into the gas previously heated by the incoming shock and the increase of the temperature after focusing is approximately given by $T/T_0 = M^4$, where M is Mach number at focusing and $T_0 \sim 300 K$ is the temperature of the ambient liquid. The observed spectrum of sonoluminescence is explained as a brehmstrahlung radiation emitted by plasma at minimum temperature $T \sim 10^5 K$.

The model reproduces nicely the time development of the bubble and sonoluminescence spectrum and explains sensitivity to the external parameters [42]. The problem is to understand how the length scales are generated and explain the jumpwise transition to sonoluminescence and the decrease of the bubble radius at sonoluminescence: ordinary hydrodynamics predicts continuous increase of the bubble radius. The length scales are the ambient radius R_0 (radius of the bubble, when gas is in pressure of 1 atm) and the minimum radius $R_s(min)$ of the shock wave determining the temperature reached in shock wave heating. Zero radius is certainly not reached since since shock front is susceptible to instabilities.

Since p-adic length scale hypothesis introduces a hierarchy of hydrodynamics with each hydrodynamics characterised by a p-adic cutoff length scale there are good hopes of achieving a better understanding of these length scales in TGD. The change in bubble size in turn could be understood as a change in the 'primary' condensation level of the bubble.

1. The bubble of air is characterized by its primary condensation level k . The minimum size of the bubble at level k must be larger than the p-adic length scale $L(k)$. This suggests that the transition to photoluminescence corresponds to the change in the primary condensation level of the air bubble. In the absence of photoluminescence the level can be assumed to be $k = 163$ with $L(163) \sim .76 \mu m$ in accordance with the fact that the minimum bubble radius is above $L(163)$. After the transition the primary condensation level of the air bubble is $k = 157$ with $L(157) \sim .07 \mu m$. In the transition the minimum radius of the bubble decreases below $L(163)$ but should not decrease below $L(157)$: this hypothesis is consistent with the experimental data [42].
2. The particles of hydrodynamics at level k have minimum size $L(k_{prev})$. For $k = 163$ one has $k_{prev} = 157$ and for $k = 157$ $k_{prev} = 151$ with $L(151) \sim 11.8 nm$. It is natural to assume that the minimum size of the particle at level k gives also the minimum radius for the spherical shock wave since hydrodynamic approximation fails below this length scale. This means that the minimum radius of the shock wave decreases from $R_s(min, 163) = L(157)$ to $R_s(min, 157) = L(151)$ in the transition to sonoluminescence. The resulting minimum radius is $11 nm$ and much smaller than the radius $.1 \mu m$ needed to explain the observed radiation if it is emitted by plasma.

A quantitative estimate goes along lines described in [42].

1. The radius of the spherical shock is given by

$$R_s = At^\alpha , \quad (5)$$

where t is the time to the moment of focusing and α depends on the equation of state (for water one has $\alpha \sim .7$).

2. The collapse rate of the adiabatically compressing bubble obeys

$$\frac{dR}{dt} = c_0 \left(\frac{2}{3\gamma} \frac{\rho_0}{\rho} \left(\frac{R_m}{R_0} \right)^3 \right)^{1/2} , \quad (6)$$

where c_0 is the sound velocity in gas, γ is the heat capacity ratio and ρ_0/ρ is the ratio of densities of the ambient gas and the liquid.

3. Assuming that the shock is moving with velocity c_0 of sound in gas, when the radius of the bubble is equal to the ambient radius R_0 one obtains from previous equations for the Mach number M and for the radius of the shock wave

$$\begin{aligned} M &= \frac{\frac{dR_s}{dt}}{c_0} = (t_0/t)^{\alpha-1} , \\ R_s &= R_0 (t/t_0)^\alpha , \\ t_0 &= \frac{\alpha R_0}{c_0} . \end{aligned} \quad (7)$$

where t_0 is the time that elapses between the moment, when the bubble radius is R_0 and the instant, when the shock would focus to zero radius in the ideal case. For $R_0 = L(167)$ (order of magnitude is this) and for $R_s(min) = L(151)$ one obtains $R_0/R_s(min) = 256$ and $M \simeq 10.8$ at the minimum shock radius.

4. The increase of the temperature immediately after the focusing is approximately given by

$$\frac{T}{T_0} \simeq M^4 = \left(\frac{R_0}{R_s} \right)^{\frac{4(1-\alpha)}{\alpha}} \simeq 1.3 \cdot 10^4 . \quad (8)$$

For $T_0 = 300 \text{ K}$ this gives $T \simeq 4 \cdot 10^6 \text{ K}$: the temperature is far below the temperature needed for fusion.

In principle the further increase of the temperature can lead to further transitions. The next transition would correspond to the transition $k = 157 \rightarrow k = 151$ with the minimum size of particle changing as $L(k_{prev}) \rightarrow L(149)$. The next transition corresponds to the transition to $k = 149$ and $L(k_{prev}) \rightarrow L(141)$. The values of the temperatures reached depend on the ratio of the ambient size R_0 of the bubble and the minimum radius of the shock wave. The fact that R_0 is expected to be of the order of $L(k_{next})$ suggests that the temperatures achieved are not sufficiently high for nuclear fusion to take place.

5.1.2 The model of sonoluminescence by Buzzacchi, del Giudice and Preparata

The coherence of the light generated in sonoluminescence looks rather mysterious from the view point of standard physics. There is very interesting paper of Buzzacchi, del Giudice and Preparata about sonoluminescence with title '*Sonoluminescence Unveiled?*' [42]. The study of this paper revealed that the physical picture behind microtubule as quantum antenna hypothesis leads to a model for sonoluminescence and that sonoluminescence could be interpreted as a direct evidence for light-like vacuum currents generating coherent photons in TGD. Needless to say, vacuum currents are a purely TGD:eish phenomenon and implied by the induced gauge field concept deriving from the hypothesis that space-time is 4-dimensional surface in certain 8-dimensional space.

The assumptions of the work of Buzzacchi, del Giudice and Preparata [42] are following.

1. The energy of the coherent radiation created in sonoluminescence results from the latent heat 0.26 eV per molecule for gas to liquid phase transition occurring at the final stage of the bubble collapse. In [42] the latent heat is used to deduce the width Γ of the energy spectrum of photons.
2. When shock wave is formed during the collapse of bubble (collapse velocity becomes supersonic), a front of layers with distance that between water molecules is formed. The average distance of molecules in tangential direction are much larger but gets smaller during the collapse of bubble. One can say that there is vapor layer looking like water in radial direction but in transversal directions the layer is much less dense. When the radius of the bubble reaches certain critical value (so that density is about 1/3 of the density of liquid phase), condensation in the transversal directions to liquid occurs. Note that this means that there is preferred direction suggesting cylindrical symmetry for the condensing regions.
3. The phase transition is assumed to occur in coherent regions with size of order $\lambda \simeq 500$ Angstroms, which is not far from the diameter of microtubules. In these regions there is a coherent planewave electromagnetic field with frequency $\omega = 2\pi/\lambda$ and the decay of this field produces the highly synchronized light flash of duration of less than 10^{-11} seconds.
4. The physical origin of the coherent regions is somewhat mysterious in standard physics but authors propose that QED is enough to explain the mystery. Authors identify the source of coherent light as resulting from the transitions between two different molecular energy states with energy difference $\Delta E \simeq 12$ eV. One could criticize this assumption as ad hoc. In any case, classical current must be defined as expectation value, which vanishes unless the two energy eigen states get mixed by interactions.

5.1.3 TGD inspired model for sonoluminescence

Consider now the TGD based modification of this model based on the same assumptions 1), 2), 3) about the origin of the coherent light as related to the liquid-gas phase transition but with different identification for the mechanism producing the coherent light. The model is based on the idea that bubble collapse might involve the sequential formation of several new space-time sheets with p-adic primes $p \simeq 2^k$, $k = 163, 157, 151, 149$ characterizing their typical sizes. The importance of the many-sheeted space-time concept was realized already in the previous rough model of the phenomenon just suggesting the identification of the basic scales of the problem in terms of the p-adic length scale hypothesis but involved no model for the generation of coherent light.

a) Light like vacuum currents generate coherent light

What is known is that the light flash emitted is *coherent* light. All frequencies are emitted simultaneously. The temporal widths of various frequencies do not depend on the nature of the gas. Thus the spectrum is certainly not genuine black body spectrum and the production mechanism must involve macroscopic quantum coherence. In TGD there indeed exists a unique mechanism leading to the generation of coherent light and is based on so called 'massless extremals' carrying light-like *vacuum currents* generating coherent light in a resonant like manner. Clearly, this mechanism predicts no dependence of the spectrum on the chemical nature of the gas in bubble. Of course, the gas can affect the spectrum by absorbing some frequencies and this indeed seems to occur. It is also known that the presence of noble gases is favorable for sonoluminescence: this is perhaps understandable from the fact that presence of noble gases (no absorption) reduces the effect of other gases by reducing their densities. In TGD inspired theory of bio-systems as macroscopic quantum systems massless extremals correspond to almost empty space-time sheets associated with microtubules and possibly also other linear bio-structures and create coherent photons (perhaps biophotons of Popp [52]).

b) Vapor-liquid phase transition in regions of microtubular size occurs

Following [42], it will be assumed shockwave formation leads to the formation of vapor layers with the mutual distance $a \simeq 3.2$ Angstroms equal to the average distance between liquid molecules and that condensation to liquid occurs when the transversal distance between the molecules of the layer becomes smaller than some critical distance $a < a_T < a_0$, where $a_0 \simeq 3.2 \times 10^{-7}$ meters is the transversal distance of the molecules when shockwave is generated. In the model of [42] $a_T \simeq \sqrt{3}a$ holds true: in TGD based model p-adic argument gives $a_T = 2a$. In TGD framework the formation of liquid phase is assumed to mean the formation of new cylindrical space-time sheets of size of order 500 Angstroms, when the transversal distance between H_2O molecules becomes critical (3 times the distance in liquid phase). At these space-time sheets water molecules are condensed into liquid phase. The length scale 500 Angstroms is suggested by [42] and in TGD framework the justification for this length scale is that it corresponds to the diameter of microtubules: these cylindrical structures could serve as templates for the formation of microtubules). Rather flat cylinders with radius equal to height are in question: of course, one can consider also cubic geometry.

c) p-Adic length scale hypothesis

p-Adic length scale hypothesis makes this picture more quantitative. Before the phase transition vapor phase is join along boundaries condensate of $k = 151$ space-time sheets glued to $k = 157$ sheets. Note that $L(151) \simeq 10^{-8}$ meters corresponds to the thickness of the cell membrane: now however the sheets are larger having size of order 500 Angstroms. Gas-to liquid phase transition is identified as a phase transition changing the value of the p-adic prime p : most naturally $k = 151 \rightarrow k = 149$. This implies $a_T = 2a_0$ rather than $\sqrt{3}a_0$ as in the model of [42]. Therefore the critical density is $\rho^* = \rho(\text{liquid})/4$ instead of $\rho(\text{liquid})/3$ of [42]. Using the relationship

$$a_T(t) = a_0 \frac{R(t)}{R_0} , \quad (9)$$

where $R_0 \simeq 4.5 \mu m$ is the radius of the bubble when contraction velocity becomes supersonic one obtains for the transversal distance a_T^* at criticality:

$$a_T^* \simeq 2a = a_0 R^* / R_0$$

giving $R^* \simeq .9 \mu m$ to be compared with $R^* \simeq .8 \mu m$ of [42].

One can estimate the thickness of the condensing shell from the requirement that the number of molecules in the shell with inner and outer radii R^* and R_0 at time t_0 is same as the number of molecules in the thin liquid shell at time when condensation to liquid has occurred. This gives for the thickness T of the liquid shell

$$T \simeq \frac{R_0^2}{R^{*2}} \frac{a^3}{a_0^3} \left(1 - \frac{R^{*3}}{R_0^3}\right) R_0 , \quad (10)$$

giving $T \simeq 10^{-7}$ meters which is *two(!)* times the size of the coherence domain as suggested by the transversal size of microtubules.

d) Topological details of the phase transition process

Consider next the topological details of the process. The transversal size of $k = 151$ sheets is (most naturally) halved in the phase transition and the join along boundaries bonds connecting $k = 151$ sheets to each other are probably split. According to the basic rules of p-adic TGD, space-time sheets with different p-adic prime p can have only wormhole contacts as stable contacts. This means that, for a p-changing phase transition to take place, the bonds connecting $k = 151$

sheets belonging to different sides of the shock front must be split, probably immediately after the formation of the shock wave. The inward flow of the newly formed $k = 149$ sheets slows down whereas the flow of $k = 151$ sheets behind them continues: the molecules condensed on them cannot however follow the flow since they collide with the liquid phase. Therefore these sheets become thus almost vacuum space-time sheets and the $k = 149$ sheets containing liquid phase topologically condense on them. At this stage the vacuum currents are generated on these almost empty $k = 151$ sheets.

e) Generation of coherent light

In the last stage of the process vacuum currents are generated on the almost empty $k = 151$ sheets and they generate the coherent light giving rise to the flash. The experience with microtubules as quantum antennae hypothesis suggests that massless extremals carrying classical light-like vacuum currents flowing in radial direction are in question. The vacuum current, possible *only* in TGD context, generates coherent photons and the flux of coherent photons from the system creates the coherent flash of photons. The frequency spectrum for the current associated with the massless extremals comes in multiples of the basic frequency π/L , L being the length of the cylinder, which is roughly equal to the thickness of H₂O layer condensing to liquid (this length is expected to have some distribution). The dependence of the vacuum current on transversal and longitudinal coordinate, which is not specified by the field equations for the vacuum extremals, in principle determines the energy spectrum. The model for the sonoluminescence should be able to predict the form of the vacuum current but this requires a model for the coupling between the parameters of the vacuum current and ordinary matter.

The model of [42] suggests that only the lowest frequency $\omega_0 = 2\pi/L$ is effectively present. The spectrum of [42] is of form

$$\begin{aligned} \frac{1}{V} \frac{dE}{d\omega} &= \frac{3\omega_0^3}{16\pi^3} |c_1|^2 |F(\omega)|^2 \frac{\omega^2}{(\omega - \omega_0)^2 + \Gamma^2/4} , \\ |c_1|^2 &\simeq 1.8 , \\ F(\omega) &= \exp(-1.4 \frac{\omega^2}{\omega_0^2}) . \end{aligned} \quad (11)$$

It is of considerable interest to verify that a spectrum, which is product of a form factor and resonance factor, results also now. The presence of the form factor $F(\omega)$ reflects the dependence of the vacuum current on transversal coordinate, which for cylindrical geometry is radial coordinate. The dependence on the transversal coordinate is left completely open by the field equations and the unknown coupling of the vacuum current with matter should determine it. The resonance factor has a purely kinematical origin: the energy spectrum for photons has form $1/(\omega - \omega_0)^2$ resulting from the fact that matrix element for photon emission involves the overlap integral $\int \exp[i(\omega - \omega_0)t] dt$ over a finite time-interval. One must take dissipation into account so that the real spectrum is proportional to $1/[(\omega - \omega_0)^2 + \Gamma^2/4]$. The resonance width is of order $\Gamma \simeq 18$ eV and in [42] it is determined by the requirement that total energy output equal to the latent heat .28 eV per molecule.

The order of magnitude for the duration of the flash can be estimated from the radial contraction velocity $dR/dt(t_0) \simeq 1.5 \times 10^3 m/s = 10^{-5}c/2$ of the bubble at the moment t_0 when the phase transition begins (according to [42]) and from the length $l \leq 500$ Angstroms, which the empty $k = 151$ sheets must travel before $k = 149$ sheets can condense on them. This gives the estimate $t(\text{flash}) \simeq 3 \times 10^{-11}$ seconds which is less than the experimental upper bound $t(\text{flash}) < 5 \times 10^{-11}$ seconds,

To summarize: sonoluminescence could provide a direct verification for the concept of massless extremal and light-like vacuum currents. Gas-liquid phase transitions could quite generally involve

the formation of massless extremals. Perhaps massless extremals of microtubular size are always present in liquid phase but carry very weak vacuum currents and bio-systems are perhaps able to amplify them. One could perhaps understand all phase transitions as formation of new space-time sheets involving p-changing phase transition.

5.2 Stirred and shaken

Japanese chemist Kazamuri Dozen and his colleagues have observed mysterious splitting of water into hydrogen and oxygen at room temperature using a simple catalyst (copper oxide in powder form) and by stirring the liquid [20]. The quicker the container is stirred the more hydrogen and oxygen are produced. Usually the dissociation occurs at temperature of about 3000 C and is driven by light: the photon density of thermal radiation has maximum at $E \sim 4T$ giving the estimate $E \sim 1.32$ eV: which gives an estimate for the energy of O–H bond possibly lowered by the presence of a catalyst. Notice that the photons in question correspond to visible light. Dömen believes that direct transformation of the kinetic energy of the liquid motion to chemical energy must take place: standard wisdom allows only the transformation *kinetic energy* \rightarrow *thermal energy* \rightarrow *chemical energy*. There is no idea about the underlying mechanism. According to [20] already 1980 analogous direct transformation of acoustic energy to chemical energy was discovered and gave rise to the field of sonochemistry. An attractive possibility is that liquid motion somehow generates coherent light which in turn drives the reaction. Similar mechanism might be at work in sonochemistry. Since sonoluminescence involves the transformation of mechanical energy into coherent light, quantum antenna hypothesis is an obvious guide line in the attempt to identify the mechanism.

The simplest TGD based mechanism explaining the anomalous splitting of hydrogen is following.

1. Stirring creates linear cylindrical vortex like structures, which are accompanied by space-time sheets carrying light like vacuum currents. The splitting to oxygen and hydrogen is driven by the coherent light emitted by the vacuum currents associated with cylindrical structures of length L . The energies for the photons of the coherent light come as multiples of $E_0 = \pi/L$ (or of $E_1 = 2\pi/L$ if periodic boundary conditions are assumed). For $E = E_0 \sim 1.32$ eV this gives the estimate $L \sim .47 \cdot 10^{-6}$ meters. This length scale is not too far from the p-adic length scale $L(163) \sim .64 \cdot 10^{-6}$ meters assuming that $L(151)$ corresponds to cell membrane thickness $L(151) \simeq 10^{-8}$ meters.
2. The rotational motion creates classical Z^0 magnetic fields (purely TGD:eish effect) realized as Z^0 magnetic flux tubes and a natural expectation is that these flux tubes are accompanied by cylindrical space-time sheets carrying light like vacuum currents. Since quarks feed their Z^0 gauge fluxes to the space-time sheets having typically twice the cell size, the naive expectation for the length of the cylindrical structures in question would be of order $L(169) \sim 5 \cdot 10^{-6}$ meters, which is however almost by one order of magnitude too large. This of course does not exclude the possibility that Z^0 magnetic flux tubes are in question. The generation of Z^0 magnetic flux tubes was suggested already many years ago to explain the observed loss of the super fluidity at much smaller critical rotational velocity than predicted by standard physics [D7].
3. A possible function of the catalyst powder is to lower the O–H bonding energy, so that it is nearer to the energy of the photons of the coherent light.

What is interesting from the point of view of consciousness theorizing is that in gel-phase vigorous streaming of intracellular liquid occurs. Furthermore, the coherent photons causing dissociation would correspond to visible light. Therefore one can wonder whether the generation of

light-like vacuum current emitting coherent biophotons [52] could be one function of the streaming. A possible test for this hypothesis is to look for an additional sink of metabolic energy inside cell.

5.3 Evidence for quantum antenna hypothesis in living systems

It is known that some monocellulars possess elementary vision based on the microtubules [35]. The emergence of the multicellulars during the Cambrian explosion was preceded by the appearance of the microtubules. If the emergence of the microtubules meant the emergence of the visual consciousness in the length scale of the cell, then the formation of the multicellulars as cell societies can be understood as a natural consequence. The length distribution of the microtubules in the rods and cones of the eye is concentrated in the region of the visible wavelengths. The coherent light in question could be identifiable as bio-photons of Popp [52].

A further piece of evidence comes from the work of Callahan about the sense of smell of insects [43]. Many insects, such as moths and ants, are known to be attracted by light, say candles and electric lamps and Callahan took as his challenge to understand what is involved. Callahan discovered that insect's olfaction is not based on chemistry but to a maser like emission of infrared light generated by various molecules such as pheromones, scent molecules and many other biomolecules. Thus insects would see rather than chemically perceive the sources of the infrared light. The sensillae of the insects serve as receiving antennas and amplify the incoming infrared radiation. Callahan also observed that the oscillation of insect antennae induce maser like emission from scent/etc. molecules by creating an oscillating emf. Thus sensory experiencing seems to involve active participation from the part of insect. The results of Callahan suggest that coherent light could be important also in our neuronal sensory experiencing.

Quite remarkably, pheromones are known to mediate sexual and social signals also in case of many mammals. For instance, certain chemical messages from female mouse can make male mouse to mate immediately while certain chemical messages from other males make him aggressive. Many mammals, for instance rodents, are known to possess vomeronasal organs, small cigar like sacks containing neurons and having length of order few millimeters [33], giving rise to an accessory olfactory system, which is known to have much more primitive structure and to work in different way than the ordinary olfactory system. It is also known that this systems bypasses cerebral cortex in rodents. There is evidence that even humans have the ability to sniff certain chemicals mediating social and sexual signals without being aware of it and there is already now an entire perfume industry based on this evidence. The chemicals giving rise to sexual attraction are probably pheromones. The fact that pheromones mediate sexual signals in case of both insects and mammals, is hardly an accident and suggests that the sensory mechanism must be the same and be based on the infrared emissions by pheromones. If the response is at neuronal level and if the cortex is not involved, one could understand why these messages are not experienced consciously. One could test this hypothesis by finding whether coherent infrared radiation at frequencies emitted by pheromones can affect the behavior of higher mammals including humans.

There is a further peculiar co-incidence: the cascade of transduction events occurring in the absorption of photon in retina is repeated in a remarkably similar way in olfactory receptor cells, which respond to odors whereas the receptor cells that respond to sound use a very different system [33]. Could this mean that also the experience of odor primarily involves the detection of (also) infrared light so that humans would not basically differ from insects or that olfactory system has evolved from the receptor neurons originally sensing infrared light? This would conform with the idea that the Kähler field generated in ear corresponds to classical Z^0 field, which does not generate coherent photons but couples with neutrinos. One must however notice that the resemblances between visual and linguistic imagery suggest that some part of ear generates cognitive representation based on coherent light and experienced by the secondary sensory organs in the thalamus.

In CASYS'2000 conference Peter Marcer reviewed the work done by him in collaboration with Russian group [45] providing experimental evidence for the hypothesis that DNA acts as receiving and sending quantum antenna. What was observed that irradiation of DNA with visible laser light induced emission of coherent light with both visible and radio frequencies. The emitted radiation was also modulated in time scale of about .01 seconds. The modulation could be due to propagation of soliton sequences propagating along Josephson junction formed by the strands of DNA or due to nonpropagating spatially constant Josephson current: both cases are mathematically equivalent with gravitational pendulum

5.4 Biefeld-Brown effect

Biefeld-Brown effect was invented as early as 1926 and is one of the oldest poorly understood electromagnetic anomalies [56, 57, 58]. The basic experiments are following.

1. Capacitor is balanced on beam balance and then charged. If the positive pole of the capacitor points upwards, the condenser moves up. If it points down the condenser moves down.
2. Capacitor is placed in free suspension such that the normal orthogonal to the plane of capacitor plates is horizontal and then charged. Capacitor is found to exhibit a horizontal thrust in the direction of the positive plate.

Thus it seems that when capacitor is provided with large charge, a force acting on capacitor in direction normal to the plane of the capacitor is observed. The motion takes place to the direction of the positively charged plate. The larger the surface area A of the capacitor, the shorter the distance d between the plates, the larger the mass M between the capacitor plates, the higher the relative dielectric constant ϵ of the dielectric, the larger the voltage V used, the larger is the size the effect. This behavior can be understood if the size of the effect is proportional to the total electric energy $E_e = \epsilon \frac{AV^2}{d}$ between capacitor plates. It is difficult to understand this effect in standard physics framework.

Consider first experiment 1) in which the normal of the capacitor plane is in vertical direction. This experiment could be understood in the general conceptual framework described in the chapter [G2]. Capacitor generates some net gravitational flux. This flux is in general fed to several space-time sheets, although most of it goes to the "standard" sheet at which the gravitational field of Earth resides. One could understand the result of the experiment a) in terms of a redistribution of these gravitational fluxes. When the positive plate points upwards/downwards the flux $\phi_{gr}(Earth)$ fed to the "standard" space-time sheet is reduced/increased. Therefore the effective weight of the capacitor decreases/increases. The dependence of $\phi_{gr}(Earth)$ on the relative orientation of the gravitational field and electric field is not surprising from TGD point of view since classical gravitational and electric fields are very closely related in TGD framework. If classical Z^0 electric force contributes to the effective gravitational force significantly, then similar mechanism in case of Z^0 electric flux could contribute significantly to the change of the effective weight of the capacitor.

It seems that this mechanism cannot explain the result of the second experiment in which capacitor moves to horizontal direction. Rather it seems that two effects must be involved. there must be some mechanism giving for the capacitor momentum in the direction of the electric field. The TGD based general mechanism of energy production relying on the generation of space-time sheets with negative time orientation and carrying negative energies could explain this aspect of Biefeld-Brown effect.

1. Suppose that the charging of capacitor involves generation of space-time sheet with negative time orientation. The energy density associated with classical fields at this space-time sheet is negative. Energy conservation requires that capacitor receives compensating energy which

in case of Biefeld-Brown effect is partially realized as kinetic energy associated with center of mass motion.

2. The classical gauge fields associated with the negative energy space-time sheet can carry also momentum and compensating momentum must be developed at the space-time sheet of the capacitor. Therefore condenser is forced to move. The momentum density of em field is proportional to the cross product $E \times B$ of the electric and magnetic fields. This momentum density gives rise to a net field momentum in the direction orthogonal to the plane of condenser plates if E and B are in directions parallel to the plates. This resembles somewhat the situation encountered in the case of Hall effect.

A working hypothesis worth of studying is that the negative energy space-time sheet associated with the capacitor corresponds to a massless extremal with E and B fields propagating from positive to negative plate (field momentum is in this direction).

1. Momentum conservation implies that the space-time sheet of the capacitor generates opposite momentum so that capacitor must move in the direction normal to the plane of the plates. What remains to be understood why the direction of motion is towards the positively charged plate. The light-likeness of 4-momentum gain together with the presence of Fourier components with single direction of wave vector means that momentum gain per energy gain is maximal. Therefore generation of negative energy "massless extremals" is optimal mechanism of propulsion. Massless extremals can have also net angular momentum since polarized Fourier components carry spin. Therefore capacitor can gain internal angular momentum in some form.
2. Assuming that the entire momentum of the classical field on negative energy space-time sheet is compensated by the momentum gain of capacitor, one obtains for the total energy gain

$$E_t = M\beta \ ,$$

where M is total mass of the capacitor and β is its velocity (the units $\hbar = 1$, $c = 1$ are used). This means quite large energy gain. For instance, for $M = .01$ kg and $\beta = 10^{-12}$, one has $E_t \sim 10^2$ Joule. The energy ΔE , which is not realized as kinetic energy, is given by

$$\Delta E = M\beta\left(1 - \frac{\beta}{2}\right) \ .$$

Obviously, only a small fraction of the energy is realized as kinetic energy of the capacitor. The ratio of the energy to thermal energy is given by

$$\frac{E_t}{E_{th}} \sim A \frac{m_p \beta}{T} \ ,$$

where A denotes atomic number. This ratio is much smaller than one for $\beta \ll T/Am_p$. In room temperature this gives $\beta \ll 10^{-11}/A$. An estimate for the magnitude of the electric field is obtained from $E_t = P$. Expressing everything in terms of integrals of energy and momentum densities, one obtains $EB \sim \rho\beta$. Since $E = B$ holds true for massless extremals, one has $B \sim \sqrt{\rho\beta}$. In condensed matter densities one has $\rho \sim Am_p/a^3$, where a is Bohr radius. This gives $B \sim \sqrt{10^5 A\beta}/a^2$. B is roughly about one \sqrt{A} Tesla for $\beta \sim 10^{-12}$. Very strong electric and magnetic fields are clearly involved.

The proposed mechanism might also make possible to understand how living systems are able to generate coherent motions.

1. The ability of bio-systems (70 per cent of water!) to generate coherent motions is complete mystery from the point of view of standard physics describing bio-system as a soup of randomly moving atoms and molecules. The generation of massless extremals provides an optimal mechanism for coherent motion. Negative energies are not absolutely essential for generating coherent motions. However, if massless extremals have positive energies, the efficiency of energy usage is however very low, approximately $\beta/2$, where β is the velocity generated: something like 10^{-8} if velocity is of order one meter per second. It could quite well be that massless extremal is created only for the period of time that motion lasts: this in accordance with the idea that classical counterparts of virtual particles are in question. Since the surplus energy generated on the material space-time sheet is partially dissipated during this time interval, this mechanism requires that metabolism feeds energy to the system to compensate this loss. Thus there is no contradiction with the general wisdom about the necessity of metabolic energy feed.
2. Brown observed that capacitors had definite effects on plants and animals. This is not surprising if TGD picture about bio-systems is correct. Coherent light is generated and this coherent light can affect the communications of neuronal society.
3. If bio-systems can generate negative energy massless extremals, a very efficient generation of metabolic energy from vacuum becomes possible. There is a lot of anecdotal evidence about the ability of yogis and mystics to survive without eating [32]. The explanation often proposed by yogis themselves [32] is that the energy of light replaces the usual sources of the metabolic energy. Standard science sceptics of course "know" and ridicule all this but, against the background of new physics predicted by TGD, I cannot avoid asking myself whether there might be some seed of truth behind these claims.

6 Quantum holography and MEs

The idea about hologrammic brain, advocated by Karl Pribram [25] and other pioneers, has several very attractive features, one of them being the robustness of the information storage due to the nonlocality of the representation. The notion of the classical hologram does not however seem to be enough as such. It is difficult to imagine how static holograms could be realized in brain. How to store simultaneously memories over entire life cycle into the same hologram at given time value is also a problem. This problem is common to all theories identifying subjective, experienced time with the geometric time of physicist: the unavoidable prediction is that brain state at given value of time must code also information about brain states at earlier times.

Peter Marcer and Walter Schempp have been advocating the notion of quantum holography as crucial for the understanding of consciousness and biocontrol [46]. Motivated by certain experimental findings Peter Gariaev and collaborators have proposed the notion wave DNA based on the idea that coherent electromagnetic fields and DNA holograms are crucial for the biocontrol [45].

Inspired by the talks of Peter Marcer in CASYS2000 and CASYS2001 conferences [47, 45], I decided to take a fresh look on how quantum holography might be realized in TGD framework, where superconducting magnetic flux tubes and massless extremals (MEs) provide an exact Bohr orbit representation of em fields and play a key role in quantum control and coordination of biomatter at atomic space-time sheets. It soon became clear that MEs are indeed tailor-made for quantum holography and teleportation in the sense as it is defined in quantum information theory [16]. Quantum holography conceptualization inspires much more detailed views about how bio-systems process information and how this information becomes conscious.

Also a profound connection with the notion of holography as it is defined in quantum gravity and string models emerges [17], and allows to deepen the vision about what the Universe predicted by quantum TGD looks like. One can also refine the formulation of quantum TGD to take

properly into account the classical non-determinism of Kähler action. In particular, the intuition that the supercanonical and superconformal symmetries of light cone boundary $\delta M_+^4 \times CP_2$, which are cosmological symmetries, generalize to approximate macroscopic symmetries acting on the light-like boundaries of the space-time sheets inside future light cone and are broken by quantum gravity, can be justified more rigorously. A rather concrete demonstration for the power of quantum holography thinking was the realization that the general ansatz for MEs should generalize followed by the observation that it indeed does so!

6.1 Quantum holography in the sense of quantum gravity theories

In string theory context quantum holography is more or less synonymous with Maldacena conjecture [18] which (very roughly) states that string theory in Anti-de-Sitter space AdS is equivalent with a conformal field theory at the boundary of AdS. In purely quantum gravitational context [17], quantum holography principle states that quantum gravitational interactions at high energy limit in AdS can be described using a topological field theory reducing to a conformal (and nongravitational) field theory defined at the *time-like* boundary of the AdS. Thus the time-like boundary plays the role of a dynamical hologram containing all information about correlation functions of $d + 1$ dimensional theory. This reduction also conforms with the fact that blackhole entropy is proportional to the horizon area rather than the volume inside horizon.

Holography principle reduces to general coordinate invariance in TGD but there are overall important delicacies involved.

1. If the action principle assigning space-time surface to a given 3-surface at light cone boundary were completely deterministic, four-dimensional general coordinate invariance would reduce the construction of the configuration geometry for the space of 3-surfaces in $M_+^4 \times CP_2$ to the construction of the geometry at the boundary of the configuration space consisting of 3-surfaces in $\delta M_+^4 \times CP_2$ (moment of big bang). Also the quantum theory would reduce to the boundary of the future light cone.
2. The classical non-determinism of Kähler action however implies that quantum holography in this strong form fails. This is indeed as it should be. The classical non-determinism (determinism in generalized sense is achieved by generalizing the notion of 3-surface by allowing unions of space-like 3-surfaces with time-like separations) is absolutely crucial for the ordinary elementary particle physics. Indeed, the quaternion-conformal symmetries responsible for the ordinary elementary particle quantum numbers act in degrees of freedom which do not contribute at all to the configuration space metric line element at all. Classical determinism would also mean that time would be lost in TGD as it is lost in GRT. Classical non-determinism is also absolutely essential for quantum consciousness and makes possible conscious experiences with contents localized into finite time interval despite the fact that quantum jumps occur between entire quantum histories.
3. Forgetting for a moment the effects of the classical non-determinism, cosmological light cone boundary $\delta M_+^4 \times CP_2$ takes the role of a quantum hologram. An important difference to AdS case is that the time-like direction of AdS boundary is replaced with light-like direction implying the metric two-dimensionality of the light cone boundary. This makes possible infinite-dimensional group of conformal symmetries in case of 4-dimensional Minkowski space. Thus the theory is much more structured than in AdS case and there is a direct connection with 2-dimensional critical systems expected on basis of quantum criticality of the TGD Universe.
4. The educated guess is that the cosmological supercanonical and superconformal symmetries associated with the light cone boundary generalize to approximate symmetries for the

sub-light cones of the cosmological light cone broken presumably only by quantum gravitational interactions. As a matter fact, not only sub-light cone boundaries but all light-like 3-dimensional boundaries inside future light cone, in particular the boundaries of massless extremals (MEs), can take the role of the cosmological light cone and define approximate supercanonical symmetries. Hence it seems that the construction of the configuration space geometry taking into account the delicacies caused by the classical non-determinism could be achieved by generalizing the construction at $\delta M_+^4 \times CP_2$ to apply also at the light-like boundaries created after big bang.

5. The generalized supercanonical symmetries commute with Poincare symmetries apart from the effects due to quantum gravitation. Therefore the states inside supercanonical representations decompose into gigantic multiplets of almost degenerate states ideal for representing biologically relevant information. MEs are especially interesting carriers of supercanonical representations.

In the following this picture is discussed in more detail.

6.2 Generalization of the solution ansatz defining massless extremals (MEs)

The solution ansatz for MEs has developed gradually to an increasingly general form and the following formulation is the most general one achieved hitherto. Rather remarkably, it rather closely resembles the solution ansatz for the CP_2 type extremals and has direct interpretation in terms of geometric optics. Equally remarkable is that the latest generalization based on the introduction of the local light cone coordinates was inspired by quantum holography principle.

The solution ansatz for MEs has developed gradually to an increasingly general form and the following formulation is the most general one achieved hitherto. Rather remarkably, it rather closely resembles the solution ansatz for the CP_2 type extremals and has direct interpretation in terms of geometric optics. Equally remarkable is that the latest generalization based on the introduction of the local light cone coordinates was inspired by quantum holography principle.

6.2.1 Local light cone coordinates

The solution involves a decomposition of M_+^4 tangent space localizing the decomposition of Minkowski space to an orthogonal direct sum $M^2 \oplus E^2$ defined by light-like wave vector and polarization vector orthogonal to it. This decomposition defines what might be called local light cone coordinates.

1. Denote by m^i the linear Minkowski coordinates of M^4 . Let (S_+, S_-, E_1, E_2) denote local coordinates of M_+^4 defining a *local* decomposition of the tangent space M^4 of M_+^4 into a direct *orthogonal* sum $M^4 = M^2 \oplus E_2$ of spaces M^2 and E^2 . This decomposition has interpretation in terms of the longitudinal and transversal degrees of freedom defined by local light-like four-velocities $v_\pm = \nabla S_\pm$ and polarization vectors $\epsilon_i = \nabla E_i$ assignable to light ray.
2. In accordance with this physical picture, S_+ and S_- define light-like curves and thus satisfy the equation:

$$(\nabla S_\pm)^2 = 0 \quad .$$

The gradients of S_\pm are obviously analogous to local light like velocities $v = (1, \bar{v})$ and $\tilde{v} = (1, -\bar{v})$. These equations are also obtained in geometric optics from Hamilton Jacobi equation by replacing photon's four-velocity with the gradient ∇S : this is consistent with the interpretation of MEs as Bohr orbits of em field.

3. With these assumptions the coordinates (S_{\pm}, E_i) define local light cone coordinates with the metric element having the form

$$ds^2 = g_{S_+ S_-} dS_+ dS_- + g_{11} dE_1^2 + g_{22} dE_2^2 .$$

Conformal transformations of M_+^4 leave the general form of this decomposition invariant. The task is to find all possible local light cone coordinates defining one-parameter families 2-surfaces defined by the condition $S_i = \text{constant}$, $i = + \text{ or } -$, dual to each other and expanding with light velocity.

6.2.2 A conformally invariant family of local light cone coordinates

The simplest solutions to the equations defining local light cone coordinates are of form $S_{\pm} = k \cdot m$ giving as a special case $S_{\pm} = m^0 \pm m^3$. For more general solutions of form

$$S_{\pm} = m^0 \pm f(m^1, m^2, m^3) , \quad (\nabla_3 f)^2 = 1 ,$$

where f is an otherwise arbitrary function, this relationship reads as

$$S_+ + S_- = 2m^0 .$$

This condition defines a natural rest frame. One can integrate f from its initial data at some two-dimensional $f = \text{constant}$ surface and solution describes curvilinear light rays emanating from this surface and orthogonal to it. The flow velocity field $\bar{v} = \nabla f$ is irrotational so that closed flow lines are not possible in a connected region of space and the condition $\bar{v}^2 = 1$ excludes also closed flow line configuration with singularity at origin such as $v = 1/\rho$ rotational flow around axis.

One can identify E^2 as a local tangent space spanned by polarization vectors and orthogonal to the flow lines of the velocity field $\bar{v} = \nabla f(m^1, m^2, m^3)$. Since the metric tensor of any 3-dimensional space allows always diagonalization in suitable coordinates, one can always find coordinates (E_1, E_2) such that (f, E_1, E_2) form orthogonal coordinates for $m^0 = \text{constant}$ hyperplane. Obviously one can select the coordinates E_1 and E_2 in infinitely many manners.

6.2.3 Closer inspection of the conditions defining local light cone coordinates

Whether the conformal transforms of the local light cone coordinates $\{S_{\pm} = m^0 \pm f(m^1, m^2, m^3), E_i\}$ define the only possible compositions $M^2 \oplus E^2$ with the required properties, remains an open question. The best that one might hope is that any function S_+ defining a family of light-like curves defines a local decomposition $M^4 = M^2 \oplus E^2$ with required properties.

1. Suppose that S_+ and S_- define light-like vector fields which are not orthogonal (proportional to each other). Suppose that the polarization vector fields $\epsilon_i = \nabla E_i$ tangential to local E^2 satisfy the conditions $\epsilon_i \cdot \nabla S_+ = 0$. One can formally integrate the functions E_i from these condition since the initial values of E_i are given at $m^0 = \text{constant}$ slice.
2. The solution to the condition $\nabla S_+ \cdot \epsilon_i = 0$ is determined only modulo the replacement

$$\epsilon_i \rightarrow \hat{\epsilon}_i = \epsilon_i + k \nabla S_+ ,$$

where k is any function. With the choice

$$k = - \frac{\nabla E_i \cdot \nabla S_-}{\nabla S_+ \cdot \nabla S_-}$$

one can satisfy also the condition $\hat{\epsilon}_i \cdot \nabla S_- = 0$.

3. The requirement that also $\hat{\epsilon}_i$ is gradient is satisfied if the integrability condition

$$k = k(S_+)$$

is satisfied: in this case $\hat{\epsilon}_i$ is obtained by a gauge transformation from ϵ_i . The integrability condition can be regarded as an additional, and obviously very strong, condition for S_- once S_+ and E_i are known.

4. The problem boils down to that of finding local momentum and polarization directions defined by the functions S_+ , S_- and E_1 and E_2 satisfying the orthogonality and integrability conditions

$$\begin{aligned} (\nabla S_+)^2 = (\nabla S_-)^2 = 0 \quad , \quad \nabla S_+ \cdot \nabla S_- \neq 0 \quad , \\ \nabla S_+ \cdot \nabla E_i = 0 \quad , \quad \frac{\nabla E_i \cdot \nabla S_-}{\nabla S_+ \cdot \nabla S_-} = k_i(S_+) \quad . \end{aligned}$$

The number of integrability conditions is 3+3 (all derivatives of k_i except the one with respect to S_+ vanish): thus it seems that there are not much hopes of finding a solution unless some discrete symmetry relating S_+ and S_- eliminates the integrability conditions altogether.

A generalization of the spatial reflection $f \rightarrow -f$ working for the separable Hamilton Jacobi function $S_{\pm} = m^0 \pm f$ ansatz could relate S_+ and S_- to each other and trivialize the integrability conditions. The symmetry transformation of M_{\pm}^4 must perform the permutation $S_+ \leftrightarrow S_-$, preserve the light-likeness property, map E^2 to E^2 , and multiply the inner products between M^2 and E^2 vectors by a mere conformal factor. This encourages the conjecture that all solutions are obtained by conformal transformations from the solutions $S_{\pm} = m^0 \pm f$.

6.2.4 General solution ansatz for MEs for given choice of local light cone coordinates

Consider now the general solution ansatz assuming that a local wave-vector-polarization decomposition of M_{\pm}^4 tangent space has been found.

1. Let $E(S_+, E_1, E_2)$ be an arbitrary function of its arguments: the gradient ∇E defines at each point of E^2 an S_+ -dependent (and thus time dependent) polarization direction orthogonal to the direction of local wave vector defined by ∇S_+ . Polarization vector depends on E^2 position only.
2. The most general MEs correspond to the solution family of the field equations having the general form

$$s^k = f^k(S_+, E) \quad ,$$

where s^k denotes CP_2 coordinates and f^k is an arbitrary function of S_+ and E . The solution represents a wave propagating with light velocity and having definite S_+ dependent polarization in the direction of ∇E . By replacing S_+ with S_- one obtains a dual solution. Field equations are satisfied because energy momentum tensor and Kähler current are light-like so that all tensor contractions involved with the field equations vanish: the orthogonality of M^2 and E^2 is essential for the light-likeness of energy momentum tensor and Kähler current.

3. The simplest solutions of the form $S_{\pm} = m^0 \pm m^3$, $(E_1, E_2) = (m^1, m^2)$ and correspond to a cylindrical MEs representing waves propagating in the direction of the cylinder axis with light velocity and having polarization which depends on point (E^1, E^2) and S_+ (and thus time).

For these solutions four-momentum is light-like: for more general solutions this cannot be the case. Polarization is in general case time dependent so that both linearly and circularly polarized waves are possible. If m^3 varies in a finite range of length L , then 'free' solution represents geometrically a cylinder of length L moving with a light velocity. Of course, ends could be also anchored to the emitting or absorbing space-time surfaces.

4. For the general solution the cylinder is replaced by a three-dimensional family of light like curves and in this case the rectilinear motion of the ends of the cylinder is replaced with a curvilinear motion with light velocity unless the ends are anchored to emitting/absorbing space-time surfaces. The non-rotational character of the velocity flow suggests that the freely moving particle like 3-surface defined by ME cannot remain in a infinite spatial volume. The most general ansatz for MEs should be useful in the intermediate and nearby regions of a radiating object whereas in the far away region radiation solution is expected to decompose to cylindrical ray like MEs for which the function $f(m^1, m^2, m^2)$ is a linear linear function of m^i .

6.3 Quantum holography in non-cosmological length scales

Quantum holography principle naturally generalizes to an approximate principle expected to hold true also in non-cosmological length and time scales.

1. The most general ansatz for MEs (inspired by the quantum holographic thinking) relies on the introduction of the notion of local light cone coordinates S_+, S_-, E_1, E_2 . The gradients ∇S_+ and ∇S_- define two light-like directions just like Hamilton Jacobi functions define the direction of propagation of wave in geometric optics. The two polarization vector fields ∇E_1 and ∇E_2 are orthogonal to the direction of propagation defined by either S_+ or S_- . Since also E_1 and E_2 can be chosen to be orthogonal, the metric of M_+^4 can be written locally as $ds^2 = g_{+-}dS_+dS_- + g_{11}dE_1^2 + g_{22}dE_2^2$. In the earlier ansatz S_+ and S_- were restricted to the variables $k \cdot m$ and $\bar{k} \cdot m$, where k and \bar{k} correspond to light-like momentum and its mirror image and m denotes linear M^4 coordinates: these MEs describe cylindrical structures with constant direction of wave propagation expected to be most important in regions faraway from the source of radiation.
2. Boundary conditions are satisfied if the 3-dimensional boundaries of MEs have one light-like direction (S_+ or S_- is constant). This means that the boundary of ME has metric dimension $d = 2$ and is characterized by an infinite-dimensional supercanonical and superconformal symmetries just like the boundary of the imbedding space $M_+^4 \times CP_2$: The boundaries are like moments for mini big bangs (in TGD based fractal cosmology big bang is actually replaced with what might be called a silent whisper amplified to not necessarily so big bang). Quantum holography would mean that effectively 2-dimensional conformal field theory at the boundary of M_+^4 region determined by ME determines what happens in the interior at QFT limit when space-time surface is not regarded as a dynamical object.
3. These observations inspire the conjecture that boundary conditions for M^4 like space-time sheets fixed by the absolute minimization of Kähler action quite generally require that space-time boundaries correspond to light-like 3-surfaces with metric dimension equal to $d = 2$. Quantum holography principle would state that the dynamics related to the metric of the configuration space, that is genuine quantum gravitation, would reduce to the boundaries of space-time sheets. The dynamics in zero modes and quaternion conformal degrees of freedom crucial for elementary particle physics would not however allow this kind of reduction. This would be consistent with the fractality which is expected to be a basic characteristic of the

quantum critical Universe predicted by TGD. The approximate supercanonical and conformal symmetries would be associated with the light-like boundaries of the space-time sheets. Supercanonical invariance would be broken only by quantum gravitational effects at the level of the configuration space by the fact that the boundaries of space-time surfaces are actually dynamical rather than fixed. The cosmological light cone boundary would be however non-dynamical and this would guarantee the exactness of the cosmological supercanonical invariance.

6.4 Supercanonical representations, quantum holography, and consciousness

Supercanonical representations are genuine quantum gravitational states since they correspond to functionals in the space of 3-surfaces (the world of worlds) and thus correspond to higher abstraction level than ordinary elementary particles. This has motivated the assignment of the so called non-geometric qualia (colors, tactile senses,...) to the quantum phase transitions associated with the supercanonical states [K3].

MEs are natural carriers of supercanonical representations obtained by multiplying ordinary physical states by configuration space Hamiltonians (functions of CP_2 coordinates and coordinates E_1, E_2 and S_+ or S_- which can obviously be arranged into irreducible representations of the color group $SU(3)$) and define an excellent candidate for a hierarchy of higher level life forms. The intuitive belief that quantum gravitation is crucial for higher level consciousness can be indeed justified in this framework: the 'worlds about worlds' aspect of higher level consciousness is what requires genuine quantum gravitational states.

The boundary of ME having one light-like direction gives rise to conformal quantum hologram representing quantum correlation functions for quantum field theory defined in the interior of ME. This 3-dimensional dynamical quantum hologram should code for conscious information about external world. This information could be determined by coherent light and gravitons scattered from the outer boundaries of other space-time sheets and could provide a quantum representation for the geometry of the boundaries of the other space-time sheets.

The standard manner to see the evolution of organism is as an initial value problem with data given at time=constant space-like section of Minkowski space. This view is definitely wrong in TGD framework, where the classical non-determinism of Kähler action is absolutely essential for the understanding of bio-systems and consciousness. Rather, one should see the problem as a boundary value problem with data given at light-like surfaces bounding MEs analogous to light cone boundary identifiable as the moment of big bang. This view conforms nicely with the active intentional aspects of the biological evolution: system can decide what it will be and life is more like a narrative with definite goals than random Brownian zigzag curve. The life cycle of the organism is specified by posing some requirements which it must satisfy in the form of boundary conditions and organism does it best to satisfy them.

6.5 Quantum holography and quantum information theory

Sokolov and collaborators [16] have proposed a model of quantum holographic teleportation in which the *classical* photocurrents from the sender to receiver take the role of a dynamical hologram. The connection with MEs is obvious.

1. MEs are carriers of classical light-like vacuum currents (one of the basic differences between TGD and Maxwell theory). This suggests that MEs could be interpreted also as *classical* holograms, which are *dynamical* as in quantum information theory. Light-like current would be like a dynamical (four-dimensional) diffraction grating. Light-like vacuum currents and vacuum Einstein tensor generate also coherent states of photons and gravitons and

MEs serve as templates for the topological condensation of photons and gravitons to the Bose-Einstein condensate of photons collinear with ME. The Bose-Einstein condensation of collinear photons (or their generalizations to 'configuration space photons') should affect the vacuum current by adding to the reference current what might be called evoked response. This condensation process could generate conscious experience and higher level qualia. Thus it would seem that MEs have a triple role as receiving and sending quantum antennae as well as classical holograms.

2. The proposal of [16] generalizes to the case of MEs provided one can devise a method of coding quantum states of photon field to the vacuum currents. The high efficiency photodetector matrix in which each pixel gives rise to a photocurrent [16], is replaced with ME or set of parallel MEs. The neural window hypothesis [H4] states that neuronal axons are accompanied by parallel MEs carrying information between sensory organs and brain and various parts of brain. This is only a less standard manner to say that ME represents classical dynamical hologram. The possibility of local light cone coordinates allows also MEs which define curved deformations of the simplest cylindrical MEs.

The concrete realization of holographic teleportation proposed in [16] brings strongly in mind the architecture of the visual pathways. Thus one can wonder whether brain is performing internal teleportation of photonic quantum states with spike patterns being directly coded to the pattern of the vacuum currents flowing along MEs. If spike patterns code the dynamical hologram, a surprisingly close relationship with Pribram's views about hologrammic brain results. Nerve pulse patterns could be seen as specifying the necessary classical aspects of the quantum teleportation (in TGD classical physics is essential part of quantum physics, rather than some effective theory).

3. Vacuum current at a 3-dimensional time-like section of ME as a function function of time defines a dynamical 3-dimensional hologram. This is consistent with the fact that our visual experience is two-dimensional: the information is always about outer boundaries of the objects of the perceptive field. The values of the vacuum current at a given point are non-deterministic which means that vacuum current is ideal for coding information. Classical data also propagate without dispersion with light velocity obeying the laws of geometric optics and MEs imply channelling so that MEs are tailor-made for classical information transfer.
4. Space-time sheets can have both positive and negative time orientations and the sign of energy depends on time orientation in TGD framework. This means that classical communication can occur both in the direction of the geometric future and past: this is essential for the classical model of the long term memories as a question communicated to the geometric past followed by answer. The dynamical nature of the holograms means that there is no need to combine 2- or 3-dimensional holograms associated with several moments of geometric time to single hologram. To remember is to perceive an object located in the geometric past. Of course, fractality might make possible temporally scaled down versions of the geometric past but the principle would remain the same.
5. Quantum hologram view suggests that the supercanonical representations at the light-like boundaries of MEs characterized by gigantic almost-degeneracies are the real carriers of biological information. According to the general theory of qualia [K3], this information would become conscious since elementary qualia would correspond to quantum jumps for which increments of the quantum numbers correspond to the quantum numbers labelling supercanonical generators in the complement of Cartan algebra. In this view superconducting magnetic flux tubes could perhaps be seen as intermediate level in the control circuitry controlled by MEs and controlling atomic level.

6. This picture leaves open whether there is a level controlling the thicknesses of the magnetic flux tubes and thus also magnetic transition frequency scales, and what this level might be. The entrainment of the endogenous frequencies to exogenous frequencies explains water memory and the effects of homeopathic remedies [54] [J2], and could make possible also endogenous NMR spectroscopy and chemical senses. The key to the puzzle might be a purely mathematical problem: how the boundary conditions at the boundaries of the magnetic flux tubes can be satisfied? It might be that the induced metric must become degenerate at the boundaries ($\sqrt{g} = 0$) implying a degeneracy of the induced metric at the boundary of the magnetic space-time sheet. This need not however mean that the M_+^4 projection of the boundary is a light-like surface: the projection could well be completely static. This supports the view that the boundaries do not carry supercanonical representations, which are associated with the imbedding space projection of the boundary rather than the boundary itself. One can imagine that ME with the same transversal section as magnetic flux tube is glued to the magnetic flux tube along this section: this kind of gluing results in a singular 4-surface analogous to the vertex region of Feynman diagram and some kind of smoothing-out procedure is needed. The smoothed-out vertex region would make possible for ME to control magnetic flux tube thickness by varying its own transversal thickness.

6.6 MEs and qualia

The properties of supercanonical representations suggests strongly that our non-geometric qualia (colors, odors,..) are associated with the supercanonical phase transitions assignable with the light-like boundaries of MEs. If so, qualia would be parameterized by the quantum number increments in the transition. The simplest qualia are in one-one correspondence with superalgebra generators which correspond to Hamiltonians labelled by spin and color quantum numbers. The notion of 'configuration space photon' defined as a ME carrying photonic Fock state and characterized by a Hamiltonian labelled by color and spin quantum numbers suggests itself. Sensory input could give rise to configuration space photons BE condensing on larger MEs. Thus photons could be genuinely colored ('qualed')!

Also magnetic quantum phase transitions give rise to qualia but it is quite possible that these qualia are not our qualia but primitive chemical qualia in much shorter length scales, say cell length scale. Magnetic qualia can however have higher level counterparts. For instance, MEs could make possible endogenous NMR and generalizations of it, and the coherent light emitted in the magnetic quantum phase transitions could BE condense on MEs and give rise to our chemical qualia.

Zero modes associated with the light-like boundaries of MEs correspond to purely geometric information about space-time surfaces and the increments of zero modes are expected to code for the purely geometric qualia (position, velocity, etc..).

Long term memories do not differ radically from ordinary qualia, the only difference being that the object of the perceptive field is in the geometric past. The light-like boundaries of ME defining the geometric correlates of selves realize this vision concretely.

6.7 Connection with the notion of wave-DNA

Peter Gariaev and his group have observed several interesting effects related to the interaction of DNA and coherent light [45]. There is evidence that the irradiation of DNA with polarized coherent light at visible wavelengths generates a radio wave response with the same polarization. Also the modulation of photon polarization by the molecules of DNA preparation is claimed to occur. There are also indications that the resulting radio waves have strange biological effects. For instance, a dramatic increase in the growth rate of potatoes irradiated by the radio waves is claimed in [45].

6.7.1 The notion of wave-DNA

These observations have led Gariaev group to propose that DNA is accompanied by a series of dynamical laser mirrors somehow storing a sequence dynamical holograms. For instance, there could be one hologram for each nucleotide pair. These effects could be indeed understood as resulting from the reading of DNA hologram by applying laser wave at a correct frequency and inducing the generation of coherent light at radio frequencies. Wave DNA hypothesis would provide a reason why for the biophotons of Popp [53].

Visible-to-radio frequency transmutation suggests that the interaction of visible photons with DNA involves the coupling of radio wave-modulated em gauge potential at visible frequencies to Dirac spinor. This coupling indeed induces a fermionic em current containing also radio frequency Fourier components in turn emitting photons at radio frequencies. The observed correlation between the polarization of the laser beam at visible frequencies with the polarization of the radio waves could be understood as reflecting the entanglement of coherent photons at these wavelengths. Note that the reduction of this entanglement in quantum jump would correspond to generation of ME self in TGD framework.

6.7.2 DNA MEs as counterparts of wave DNA

In standard physics framework a possible objection against this scenario is that the wavelengths of the visible photons are almost one micrometer and much longer than the transversal DNA length scale. In TGD framework the laser mirrors associated with DNA could be single or double sheeted MEs orthogonal to DNA. Since these MEs represent space-time sheets, their length could quite well be measured in multiples of visible light wavelength. Note that the thickness of chromosomes is of order $E - 7$ meters, not far from visible wavelengths.

DNA MEs would be analogous the ELF MEs assumed to drift along cortex and scan and control the state of various brain regions [K3]. The variation of the magnetic field along DNA would induce the drifting force. Of course, DNA MEs need not drift. DNA MEs could be like pages of a book [45] with each page storing its own independent dynamical hologram realized as a light-like classical vacuum current. The local polarization direction of the classical em field inside ME varies continuously and depends on the light-like coordinate S_+ (on S_- for dual ME). This means that a dynamical orientation field in plane can be represented by single ME. The local polarization of the coherent light is indeed proposed to be carrier of bio-information in [45].

DNA ME would describe a sawtooth like zigzag curve in space-time. At the turning points it would be glued to other space-time sheets representing matter: vacuum current and information content would be determined by what happens at these two-dimensional regions.

Quantum gravitational hologram principle says that the quantum information about the development of the organism is stored by the supercanonical representations which can be thought of as localized to the light-like boundaries of MEs. The immense almost degeneracies of supercanonical states would indeed mean huge information storage capacities and almost infinite adaptivity already at DNA level. The information stored by the genome would be only a tiny part of the information and comparable to the read-only-memory burnt into the hardware of computer.

6.7.3 DNA solitons and phantom DNA effect as evidence for superconducting DNA space-time sheets

According to [45] there is also evidence for electro-acoustic soliton sequences propagating along DNA and inducing a local opening of DNA double strand. In TGD framework these soliton sequences could be seen as scaled-down versions of nerve pulse patterns. They would be induced from the Sine-Gordon soliton sequences (the simplest soliton sequences are equivalent with a rotational motion of gravitational pendulum propagating with sub- or superluminal phase velocity). The

solitons would be associated with electron (most naturally) Josephson currents running between DNA strand space-time sheets connected by Josephson junctions idealizable with a continuous Josephson junction. In accordance with homeostasis as an ionic flow equilibrium idea, stochastic resonance could induce acoustic solitons at atomic space-time sheets involving the local opening of DNA. MEs, in particular DNA MEs, could serve as ultimate quantum controllers inducing electron super currents at various space-time sheets associated with DNA and thus affecting the ionic flow equilibrium at the atomic space-time sheets.

MEs and superconducting magnetic flux tubes associated with DNAs could also explain the phantom DNA effect [48]. Some of the magnetic flux tubes would be left to chamber after removal of DNA and provide a representation of quantum control level of DNA. The intensity of the radiation generated in phantom DNA effect would give an estimate for the density of ions at the superconducting magnetic flux tube space-time sheets. Water memory would be also based on similar structures.

6.8 p-Adic cognition and the existence of double-sheeted MEs and wormhole magnetic fields

By the inherent non-determinism of the p-adic differential equations, p-adic space-time sheets represent a natural candidate for the geometric correlates of imagination. The transformation of thought into action corresponds in p-adic-to-real phase transition in this conceptual framework. TGD predicts that space-time sheets with negative time orientation carry negative energies. Thus conservation laws allow p-adic–real transitions for pairs of space-time sheets having opposite energies and momenta. Presumably it is also easier to satisfy the boundary conditions at the ends of the double sheeted structures of finite length.

In particular, the MEs associated with cognition should be double sheeted structures with vanishing net energy. In this case the classical fields could be precise time reversals (phase conjugates) of each other. The analogy with DNA double sheet is obvious. Also DNA MEs could be double sheeted and represent cognitive structures at molecular level.

The double sheeted MEs could represent dynamical holograms associated with long term memory: negative energy ME would represent a question communicated to the geometric past and positive energy ME would represent the answer of self woken up in geometric past and by self-organization generating the answer. To remember would be to make a quantum jump generating this kind of double sheeted ME. Subjectively this would be completely instantaneous process and could involve the generation of entanglement between self geometrically now and self in in geometric past and its reduction in quantum jump. Communication/long term memory could be seen as a four-dimensional quantum measurement.

MEs are not the only double-sheeted cognitive structures one can imagine. The so called wormhole magnetic fields are formed by pairs of space-time sheets carrying opposite magnetic fields and are analogous to double sheeted MEs and DNA double strands. Magnetic space-time sheets are connected by extremely tiny wormhole contacts. The wormhole throats of the wormhole contact carry opposite classical em charges and rotating wormhole contacts induce opposite magnetic fields at two flux tube space-time sheets. Bio-systems might be able to generate both double sheeted MEs and wormhole magnetic fields at will by p-adic-real phase transitions since energetic constraints would not pose any restrictions. The p-adic versions of wormhole magnetic fields would serve as cognitive representations for Earth's magnetic field and the magnetic circulation associated with body could involve also wormhole magnetic fields. Quite generally, TGD Universe might be mimicking itself by constructing double-sheeted zero energy copies of single sheeted structures representing matter. This process might be an essential aspect of water memory and homeopathy [J5].

6.9 Connection with quantum holography involved with NMR

A highly abstract and refined form of quantum holography appears also in the mathematical formulation of NMR spectroscopy by Walter Schempp [19]. In this formulation coherent photon states are identified as states in the irreducible unitary representations of the Heisenberg group generated by a bosonic oscillator operator, its conjugate and identity operator. The vision about homeostasis as many-sheeted ionic flow equilibrium encourages the idea that endogenous NMR and its generalizations to arbitrary magnetic transitions make possible chemical senses.

1. The ultracold magnetic flux tubes carrying superconducting ions and molecules are ideal for NMR. The contact with ME perturbs the magnetic field inside a magnetic flux tube by superposing to it a rotating transversal component. The perturbations of Earth's magnetic field have frequencies in EEG range (by purely geometric arguments). More generally, the inverse of the geometric size scale for the flux tubes defines the minimum frequency for the perturbations. The temperature is extremely low so that the ratio $N_+/N_- = \exp(-\Delta E/kT)$ of ions with different spin directions in magnetic field associated with magnetic flux tube (of say Earth's magnetic field) deviates strongly from unity. This makes possible to avoid the immense reduction of NMR signal intensity (proportional to $\exp(-\Delta E/kT) - 1$) occurring in the ordinary NMR spectroscopy. If magnetic quantum transitions occur coherently for the superconducting BE condensate, the rate for quantum transition is proportional to the square of the number of superconducting ions and this gives an additional amplification factor. The rate for the BE condensation of the coherent light generated in the magnetic quantum phase transition is also amplified by the presence of the collinear BE condensed photons at MEs acting as receiving quantum antennae.
2. If the magnetic flux tube thickness varies along the flux tube, the strength of the magnetic field and magnetic transition frequency scale vary too. In this kind of situation the interaction produces NMR spectroscopy for superconducting ions at superconducting magnetic flux tubes. The resonance frequency spectrum place-code various ionic concentrations in the sense that each ion would give to a conscious resonance located at a particular position of the magnetic flux tube. This would be one example about the coding of abstract geometric data (now ionic concentrations) to magnetic field strength to position along magnetic flux tube.
3. Magnetic phase transitions could represent some kind of primitive chemical qualia, much more primitive than the qualia associated with the transitions of genuinely quantum gravitational super-canonical states at MEs (non-trivial functionals in the world of worlds, space of 3-surfaces). The magnetic qualia need not be conscious to us since the characteristic length scale given by the magnetic length is cell length scale in the case of endogenous magnetic field $B_{end} = .2$ Gauss which has same order of magnitude as Earth's magnetic field with nominal value $B_E = .5$ Gauss. The value of B_{end} comes from the explanation for the effects of ELF em fields on vertebrate in terms of cyclotron transitions. The BE condensation of the coherent light emitted in the magnetic phase transitions on MEs could however lead to higher level variants of the chemical qualia.
4. In the many-sheeted ionic flow equilibrium superconducting ion densities are related to the ionic densities at the atomic space-time sheets. This means that the information about ionic densities at the atomic space-time sheets could be coded into vacuum currents and MEs could contain holographic representation of the biochemical state of the organism. The possibility of Z^0 magnetic flux tubes would also make possible to code information about the densities of neural atoms and molecules to Z^0 MEs. Unless massless exotic variants of Z^0 bosons are possible, higher level variants of Z^0 type chemical qualia are not possible.

6.10 Summary

It seems that the vision about quantum control and experiencing is becoming rather concrete and beautiful overall pattern is emerging. At least following anomalous from standard physics point of view pieces fit together.

1. The paradoxical results related to the ionic channels and pumps [36] give very strong support for ionic flow equilibrium in many-sheeted space-time.
2. Water memory and homeopathy fits with this picture. The results of Cyril Smith [54] about homeopathic effects at various frequencies as well as the findings of Benveniste [39, 40] and recent findings of a collaboration involving four laboratories about water memory [51] support this view. The observed effects of ELF em fields on biomatter [44] are also consistent with the assumption that ELF em fields affect directly the control circuitry at superconducting space-time sheets. The biophotons of Popp [53] should also be mentioned as also the puzzles related to the interaction of biomolecules (hormones, second messengers) with cell nucleus (pleiotropy) suggesting that non-chemical communication mechanisms are involved.
3. A common answer to the questions 'Why liquid crystals?', 'Why electrets?', 'Why DC currents of Becker [38]?', 'Why collagen networks?', is that otherwise the many-sheeted control circuitry is not possible.
4. The healing of bone fractures by electric currents discovered by Becker [38] could also be a manner to re-establish super current patterns leading to many-sheeted ionic flow equilibrium and healing. Note the possible applications in nanotechnology: the generation of an appropriate supra current pattern could force ionic flow equilibrium leading to the healing of fractures in electromechanical systems.
5. The recent empirical findings [31] provide strong support for the belief that NDE experiences and thus consciousness are possible even when brain is clinically dead and it is interesting to ponder what this might mean in TGD framework. Both MEs and magnetic flux tube structures form a fractal hierarchy. Rather interestingly, the magnetic fields associated with eyes have cyclotron time scale varying from hour to year roughly. This is necessary for higher abstraction levels of consciousness since they involve subjective memories in time scale much longer than the .1 seconds of sensory experience. Flux tubes are of thickness of order millimeter in this case: the size scale of iris!

Very rough estimates suggest that the magnetic crystals contained by brain (say those in pineal gland) could generate fields in the range $10^{-2} - 1$ fT, flux tube thickness would be in the range 1 – 10 cm and cyclotron time scale would vary from years to hundred years (life summary during NDE). This provokes several questions. Could these magnetic flux tubes connect near relatives, like umbilic cords (meeting of relatives during NDEs)? Could they give rise to tunnel experience involved with NDEs: magnetic flux tube self experiencing fractal hierarchy of sub-flux tubes as mental images (my own 'tunnel like' visual experiences after sitting and writing resemble incompressible fluid flow, $\nabla \cdot B = 0$)? Is this magnetic consciousness what remains after physical death: are our bodies sensory and motor organs of huge fractal magnetic circuitry ('light being') living in symbiosis with MEs?

7 Appendix: A model for the topological condensation of coherent vapor phase photons

In ordinary QED classical gauge fields can have only ordinary charged particles as their sources. In TGD genuine vacuum currents are possible. The coupling of the quantum field to the classical

em field with a non-vanishing vacuum source implies the generation of a coherent state of photons such that each Fourier component present in the classical gauge current gives rise to an eigen state of the corresponding photonic annihilation operator. In case of light-like vacuum currents allowed by TGD, the coherent state is generated in resonant-like manner so that light-like vacuum current acts as an ideal quantum antenna.

If one introduces a second space-time sheet, which contains BE condensate of photons for some modes of the photon field, a stimulated topological condensation of both coherent vapor phase photons and transfer of coherent condensed photons from other space-time sheets to this space-time sheet occurs. This effect makes possible the the action of the second space-time sheet as an optimal receiving antenna. In the following calculation the consideration is restricted to the stimulated condensation of vapor phase photons.

In biological context microtubules could server both as senders and receivers of coherent photons. According to the proposed identification of coherent photons as the quantum correlate of vision, the microtubules contain BE condensate of photons in some some modes would have the ability to see in some primitive manner.

7.1 The action

The simplest model for the situation is based on Maxwell action for electromagnetic field regarded as an induced field obtained from superposition of the classical emf in CP_2 degrees of freedom and second quantized free emf in $M_+^4 \times CP_2$ having only M^4 components and depending on M_+^4 coordinates only and having decomposition into vapor phase and condensate parts ($\hbar = 1$ and $c = 1$ will defined the units used in the following).

$$\begin{aligned}
F_{\mu\nu} &= F_{\mu\nu}(cl) + F_{\mu\nu}(qu) , \\
F_{\mu\nu}(cl) &= F_{kl}(cl)\partial_\mu s^k \partial_\nu s^l , \\
F_{\mu\nu}(qu) &= F_{kl}(qu)\partial_\mu m^k \partial_\nu m^l , \\
F_{kl}(qu) &= \partial_l A_k(qu) - \partial_k A_l(qu) , \\
A_k(qu) &= A_k(qu, v) + A_k(qu, c) .
\end{aligned} \tag{12}$$

$F_{kl}(qu)$ satisfies empty space Maxwell equations. m^k and s^k refer to M_+^4 and CP_2 coordinates and v and c refer to vapor phase and condensate.

Maxwell action density can be transformed to a sum of a total divergence reducing to mere boundary term, to be neglected, plus free part and two interaction terms in the following manner:

$$\begin{aligned}
\frac{L}{\sqrt{g}} &= \sum_i L(\text{free}, i) + L_1(\text{int}) + L_2(\text{int}) , \\
L(\text{free}, i) &= \frac{1}{4} F_{\mu\nu}(qu, i) F^{\mu\nu}(qu, i) , \quad i = c, v . \\
L_1(\text{int}) &= \frac{1}{2} j^\mu(cl) \sum_i A_\mu(qu, i) , \\
L_2(\text{int}) &= \frac{1}{2} \sum_i J^\mu(qu, i) A_{m\mu}(cl) , \\
J^\mu(qu, i) &= F_k^\mu(i) M_\nu^{k\nu} + F_k^\nu(i) M_\nu^{k\mu} , \quad i = c, v \\
M_{\alpha\beta}^k &= D_\beta \partial_\alpha m^k .
\end{aligned} \tag{13}$$

$L(\text{free}, i)$ denotes the free action for the classical emf and vapor phase and condensed quantum emfs and defines photon propagators. Standard propagator is obtained, when Minkowski coordinates are used for space-time surface.

$L_1(\text{int})$ corresponds to the action of the vapor phase and condensed quantum emf with the vacuum current and leads to generation of coherent state of photons both in vapor phase and condensate.

$L_2(\text{int})$ is non-vanishing only, when the M_+^4 part of the second fundamental form $M_{\alpha\beta}^k$ for 4-surface is non-vanishing: in this case the em current associated with $A_{mu}(qu)$ is non-vanishing despite the fact that it vanishes for $A_k(qu)$! This term describes the external curvature of the 4-surface as opposed to the internal curvature described by the curvature tensor. In general case, the external curvature can be large even when the gravitational field vanishes. It must be however emphasized that this term is proportional to the metric of CP_2 and, in case of the massless extremals, this term is significant only if the dependence of CP_2 coordinates on the transversal coordinates of M_+^4 is strong: this in turn requires huge value for the light-like Einstein tensor. This term will be neglected in the sequel.

The representation

$$\begin{aligned} A_+(k, \lambda) &= \sqrt{\frac{2\pi}{\omega_k}} a^\dagger(k, \lambda) , \\ [a(k_1, \lambda_1), a^\dagger(k_2, \lambda_2)] &= \delta^3(k_1 - k_2) \delta_{\lambda_1, \lambda_2} , \end{aligned} \quad (14)$$

for which the density of states factor for photon states is $dN = d^3k$, will be used in the sequel.

7.2 Coherent state is generated in resonant-like manner for light-like vacuum currents

The presence of the vacuum current leads to the generation of coherent state of photons both in vapor phase and condensate. Coherent states are eigen states of the photonic annihilation operators and in the estimates for the rate of topological condensation, one in a good approximation one can replace $A_\mu(qu, i)$, $i = \text{cond}, \text{vap}$, with the classical photon field $A_\mu(\text{coh}, i)$ having classical vacuum current as its source and serving as order parameters for the coherent state. The Fourier component of a vector potential describing the eigenvalue of the annihilation operator part of the photon field is for given momentum k and polarization direction λ given by

$$\begin{aligned} A^\mu(\text{coh}, v|\lambda, k) &= \sum_n c(k, k_n) \frac{\lambda_\mu J^\mu(k_n) \lambda^\mu}{k_n^2} , \\ \exp(-ik \cdot m) &= \sum_n c(k, k_n) \exp(-ik_n \cdot m) . \end{aligned} \quad (15)$$

$c(k, k_n)$ is the Fourier component of the planewave $\exp(-ik \cdot m)$ expressed using discrete planewave basis for the space-time sheet containing the vacuum current. m denotes Minkowski coordinates.

If the classical vacuum current is associated with a 'massless extremal', em current is light-like and this implies resonance for those frequencies for which photon wave vector corresponds to the wave vectors appearing in the vacuum current. The resonance is smoothed out by the finite spatial size of the space-time sheet containing the light-like vacuum current. At the limit of an infinitely large spatial size for the space-time sheet, one obtains infinitely large amplitude since one has $k_n^2 = k^2 = 0$ at this limit.

7.3 Stimulated topological condensation

The presence of the coherent state of photons implies the possibility of the topological condensation of photons. If the device contains $N(k, \lambda)$ photons in the state (k, λ) , stimulated topological condensation, completely analogous to the stimulated emission, occurs and the condensation rate is proportional to $(N(k, \lambda) + 1)^2$.

Assume that there exists a coherent state generated by quantum antenna of possibly astrophysical dimension and associate label '1' with this space-time sheet. Assume also a second space-time sheet and associate with it label '2'. In the lowest order the matrix element for the topological condensation of single photon can be obtained as the matrix element of the creation operator part of the interaction term of the action

$$\begin{aligned}
iS_+ &= \frac{i}{2} \int_{V_2} dV_2 j_\perp^\mu(\text{coh}, 1) A_{\mu,+}(\text{cond}, 2) \\
&= \frac{i}{2} \sum_{\lambda_2} \int d^3 k_2 X(k_2, \lambda_2) a^\dagger(k_2, \lambda_2) , \\
X(k_2, \lambda_2) &= \sqrt{\frac{2\pi}{\omega_{k_2}}} \sum_{\lambda_1} \int d^3 k_1 Y(k_1, \lambda_1, k_2, \lambda_2) , \\
Y(k_1, \lambda_1, k_2, \lambda_2) &= j(\text{coh}, 1|k_1, \lambda_1) c(k_1, k_2) e_{\lambda_1} \cdot e_{\lambda_2} , \\
c(k_1, k_2) &= \int_{V_2} dV_2 \exp[i(k_1 - k_2) \cdot m] ,
\end{aligned} \tag{16}$$

between the initial and final states. $j^\mu(\text{coh}, 1)$ is just the transversal part of the classical vacuum current creating the coherent state. The latter expression is obtained by using Fourier expansions for j and A_+ (, which denotes the creation operator part of the free photon field projected to the space-time surface representing the device: Minkowski coordinates are used for both source regions and device).

In case that the region V_1 is box of length L in the direction of the vacuum current, the explicit calculation, writing the light-like vacuum current as $j^\mu = Jp^\mu$, $p^0 = p^z = 1$, leads to the following expression for the Fourier component $j(\text{coh}, 1|k_1, \lambda_1)$:

$$\begin{aligned}
j(\text{coh}, 1|k, \omega_k, \lambda) &= j^\mu(\text{coh}, 1|k, \omega_k) e_\mu^\lambda , \\
&= \sum_n \frac{\exp(ik_z L) - 1}{ik_z} J(\omega_n, k_T) p \cdot e^\lambda \delta(k^0 - \omega_n) , \\
\omega_n &= \frac{n\pi}{L} .
\end{aligned} \tag{17}$$

Delta-function expresses the fact that only discrete frequencies are allowed for the vacuum current and one can write the condensation amplitude as a sum $iS_+ = i \sum_n iS_{+,n}$ over the allowed frequencies ω_n . k_T refers to the transversal part of the wave vector orthogonal to the light-like vacuum current.

From this expression one can deduce the probability for the topological condensation of photon (k, λ) to a state containing $N(k, \lambda)$ photons as

$$|S(k, \lambda)|^2 = \left| \sum_n S_{+,n} \right|^2 (N(k, \lambda) + 1)^2 , \tag{18}$$

Clearly, $(N(k, \lambda) + 1)^2$ factor corresponds to the induced condensation. By a standard trick one can eliminate the square of the delta-function by replacing the condensation probability with condensation rate $R(k, \lambda)$ obtained by dividing condensation probability with $T \rightarrow \infty$ eliminating one deltafunction. Furthermore, one can calculate the transition rate to a set of final states by multiplying the expression thus obtained with the density of states factor $dN = d^3 k$, which after the elimination of the second delta function effectively reduces to $\omega_n^2 d\Omega$. In this manner one obtains for the differential condensation rate a rather neat expression in terms of the vacuum current

$$\begin{aligned}
\frac{dR(k,\lambda,n)}{d\Omega} &= \frac{\pi}{2}\omega_n L^2 |M(k,\lambda)|^2 (N(k,\lambda) + 1)^2, \\
M(k,\lambda) &= i \sum_{\lambda_1} \int d^3 k_1 J(\omega_n, k_T^1) c(k^1, k) X(k_1, \lambda_1), \\
X(k_1, \lambda_1) &= \frac{\exp(ik_z^1 L) - 1}{ik_z^1 L} p \cdot e_{\lambda_1} e_{\lambda_1} \cdot e_{\lambda}.
\end{aligned}
\tag{19}$$

From this expression it is clear that resonance indeed occurs and at the limit $L \rightarrow \infty$ the rate for condensation diverges as L^2 . In this expression the overlap integral $c(k_1, k_2)$ carries information about the geometry of the space-time sheet associated with the 'device' whereas $J(\omega_n, k_T)$ characterizes the vacuum current and the remaining factor X is a purely 'kinematic' factor.

References

Online books about TGD

- [1] M. Pitkänen (2006), *Topological Geometroynamics: Overview*.
<http://www.helsinki.fi/~matpitka/tgdview/tgdview.html>.
- [2] M. Pitkänen (2006), *Quantum Physics as Infinite-Dimensional Geometry*.
<http://www.helsinki.fi/~matpitka/tgdgeom/tgdgeom.html>.
- [3] M. Pitkänen (2006), *Physics in Many-Sheeted Space-Time*.
<http://www.helsinki.fi/~matpitka/tgdclass/tgdclass.html>.
- [4] M. Pitkänen (2006), *Quantum TGD*.
<http://www.helsinki.fi/~matpitka/tgdquant/tgdquant.html>.
- [5] M. Pitkänen (2006), *TGD as a Generalized Number Theory*.
<http://www.helsinki.fi/~matpitka/tgdnumber/tgdnumber.html>.
- [6] M. Pitkänen (2006), *p-Adic length Scale Hypothesis and Dark Matter Hierarchy*.
<http://www.helsinki.fi/~matpitka/paddark/paddark.html>.
- [7] M. Pitkänen (2006), *TGD and Fringe Physics*.
<http://www.helsinki.fi/~matpitka/freenergy/freenergy.html>.

Online books about TGD inspired theory of consciousness and quantum biology

- [8] M. Pitkänen (2006), *Bio-Systems as Self-Organizing Quantum Systems*.
<http://www.helsinki.fi/~matpitka/bioselforg/bioselforg.html>.
- [9] M. Pitkänen (2006), *Quantum Hardware of Living Matter*.
<http://www.helsinki.fi/~matpitka/bioware/bioware.html>.
- [10] M. Pitkänen (2006), *TGD Inspired Theory of Consciousness*.
<http://www.helsinki.fi/~matpitka/tgdconsc/tgdconsc.html>.
- [11] M. Pitkänen (2006), *Mathematical Aspects of Consciousness Theory*.
<http://www.helsinki.fi/~matpitka/genememe/genememe.html>.

- [12] M. Pitkänen (2006), *TGD and EEG*.
<http://www.helsinki.fi/~matpitka/tgdeeg/tgdeeg/tgdeeg.html>.
- [13] M. Pitkänen (2006), *Bio-Systems as Conscious Holograms*.
<http://www.helsinki.fi/~matpitka/hologram/hologram.html>.
- [14] M. Pitkänen (2006), *Magnetospheric Consciousness*.
<http://www.helsinki.fi/~matpitka/magnconsc/magnconsc.html>.
- [15] M. Pitkänen (2006), *Mathematical Aspects of Consciousness Theory*.
<http://www.helsinki.fi/~matpitka/magnconsc/mathconsc.html>.

References to the chapters of books

- [A1] The chapter *An Overview about the Evolution of Quantum TGD* of [1].
<http://www.helsinki.fi/~matpitka/tgdview/tgdview.html#tgdevo>.
- [C14]
- [C6] The chapter *Was von Neumann Right After All* of [4].
<http://www.helsinki.fi/~matpitka/tgdquant/tgdquant.html#vNeumann>.
- [D6] The chapter *TGD and Astrophysics* of [3].
<http://www.helsinki.fi/~matpitka/tgdclass/tgdclass.html#astro>.
- [D7] The chapter *Macroscopic Quantum Phenomena and CP_2 Geometry* of [3].
<http://www.helsinki.fi/~matpitka/tgdclass/tgdclass.html#super>.
- [F3] The chapter *p-Adic Particle Massivation: Hadron Masses* of [6].
<http://www.helsinki.fi/~matpitka/paddark/paddark.html#padmass2>.
- [F6] The chapter *Topological Condensation and Evaporation* of [6].
<http://www.helsinki.fi/~matpitka/paddark/paddark.html#padaelem>.
- [F8] The chapter *TGD and Nuclear Physics* of [6].
<http://www.helsinki.fi/~matpitka/paddark/paddark.html#padnucl>.
- [F9] The chapter *Dark Nuclear Physics and Living Matter* of [6].
<http://www.helsinki.fi/~matpitka/paddark/paddark.html#exonuclear>.
- [G2] The chapter *The Notion of Free Energy and Many-Sheeted Space-Time Concept* of [7].
<http://www.helsinki.fi/~matpitka/freenergy/freenergy.html#freenergy>.
- [G3] The chapter *Did Tesla Discover the Mechanism Changing the Arrow of Time?* of [7].
<http://www.helsinki.fi/~matpitka/freenergy/freenergy.html#tesla>.
- [H1] The chapter *Matter, Mind, Quantum* of [10].
<http://www.helsinki.fi/~matpitka/tgdconsc/tgdconsc.html#conscic>.
- [H3] The chapter *Self and Binding* of [10].
<http://www.helsinki.fi/~matpitka/tgdconsc/tgdconsc.html#selfbindc>.
- [H4] The chapter *Quantum Model for Sensory Representations* of [10].
<http://www.helsinki.fi/~matpitka/tgdconsc/tgdconsc.html#expc>.

- [H7] The chapter *Conscious Information and Intelligence* of [10].
<http://www.helsinki.fi/~matpitka/tgdconsc/tgdconsc.html#intsysc>.
- [I1] The chapter *Quantum Theory of Self-Organization* of [8].
<http://www.helsinki.fi/~matpitka/bioselforg/bioselforg.html#selforgac>.
- [I4] The chapter *Quantum Control and Coordination in Bio-systems: Part I* of [8].
<http://www.helsinki.fi/~matpitka/bioselforg/bioselforg.html#qcococI>.
- [I5] The chapter *Quantum Control and Coordination in Bio-Systems: Part II* of [8].
<http://www.helsinki.fi/~matpitka/bioselforg/bioselforg.html#qcococII>.
- [J1] The chapter *Bio-Systems as Super-Conductors: part I* of [9].
<http://www.helsinki.fi/~matpitka/bioware/bioware.html#superc1>.
- [J2] The chapter *Bio-Systems as Super-Conductors: part II* of [9].
<http://www.helsinki.fi/~matpitka/bioware/bioware.html#superc2>.
- [J4] The chapter *Quantum Antenna Hypothesis* of [9].
<http://www.helsinki.fi/~matpitka/bioware/bioware.html#tubuc>.
- [J5] The chapter *Wormhole Magnetic Fields* of [9].
<http://www.helsinki.fi/~matpitka/bioware/bioware.html#wormc>.
- [J6] The chapter *Coherent Dark Matter and Bio-Systems as Macroscopic Quantum Systems* of [9].
<http://www.helsinki.fi/~matpitka/bioware/bioware.html#darkbio>.
- [K1] The chapter *Time, Spacetime and Consciousness* of [13].
<http://www.helsinki.fi/~matpitka/hologram/hologram.html#time>.
- [K3] The chapter *General Theory of Qualia* of [13].
<http://www.helsinki.fi/~matpitka/hologram/hologram.html#qualia>.
- [K4] The chapter *Bio-Systems as Conscious Holograms* of [13].
<http://www.helsinki.fi/~matpitka/hologram/hologram.html#hologram>.
- [K5] The chapter *Homeopathy in Many-Sheeted Space-Time* of [13].
<http://www.helsinki.fi/~matpitka/hologram/hologram.html#homeoc>.
- [K6] The chapter *Macroscopic Quantum Coherence and Quantum Metabolism as Different Sides of the Same Coin* of [13].
<http://www.helsinki.fi/~matpitka/hologram/hologram.html#metab>.
- [L1] The chapter *Genes and Memes* of [11].
<http://www.helsinki.fi/~matpitka/genememe/genememe.html#genememec>.
- [L2] The chapter *Many-Sheeted DNA* of [11].
<http://www.helsinki.fi/~matpitka/genememe/genememe.html#genecodec>.
- [L4] The chapter *Pre-Biotic Evolution in Many-Sheeted Space-Time* of [11].
<http://www.helsinki.fi/~matpitka/genememe/genememe.html#prebio>.
- [M2] The chapter *Quantum Model for Nerve Pulse* of [12].
<http://www.helsinki.fi/~matpitka/tgdeeg/tgdeeg/tgdeeg.html#pulse>.
- [M3] The chapter *Dark Matter Hierarchy and Hierarchy of EEGs* of [12].
<http://www.helsinki.fi/~matpitka/tgdeeg/tgdeeg/tgdeeg.html#eegdark>. T

- [M4] The chapter *Quantum Model for EEG: Part I* of [12].
<http://www.helsinki.fi/~matpitka/tgdeeg/tgdeeg/tgdeeg.html#eegI>.
- [M5] The chapter *Quantum Model of EEG: Part II* of [12].
<http://www.helsinki.fi/~matpitka/tgdeeg/tgdeeg/tgdeeg.html#eegII>.

Theoretical physics

- [16] I. V. Sokolov *et al*(2001), *Quantum holographic teleportation of light fields*, arXiv:quantph/0007026v1 .
- [17] S. de Haro Olle (2001), *Quantum Gravity and the Holographic Principle*, thesis. arXiv:hep-th/0107032v1 .
- [18] J. M. Maldacena (1998), *The Large N Limit of Superconformal Field Theories and Super Gravity*, Adv. Theor. Math. Phys. 2, 231-252. hep-th/9802150 .
- [19] W. Schempp (1998), *Magnetic Resonance Imaging, Mathematical Foundations and Applications*, John Wiley, New York.
- [20] L. Guterman (1998), *Stirred and Shaken*, New Scientist, No 2160, Nov. 14.

Biology and neuroscience

- [21] H. Baars, (199?), *The neural basis of conscious experience*.
<http://www.wrightinst.edu/faculty/baars/book/text/CTC-ch03.htm>.
- [22] E.R. Kandel, J.H. Schwartz, T. M. Jessel (1991), *Principles of neural science*, Prentice-Hall International Inc. .
- [23] Ann Gibbons (1998), *Solving the Brain's Energy Crisis*, Science, Vol. 280, 29 May 1998.
- [24] K. Matter-Walstra (1999), *Unasked questions*.
<http://www.datacomm.ch/kmatter/unaskq.htm> .
- [25] Pribram, K. H. (1969), *The Neurophysiology of Remembering*. Scientific American, January 1969.
 K. H. Probram (1991), *Brain and Perception; Holonomy and Structure in Figural Processing*, Lawrence Erlbaum Associates, New Jersey.
- [26] C. B. Pert (1997), *Molecules of Emotion*, Simon & Schuster Inc..
- [27] R. Penrose(1994) *Shadows of the Mind*, Vintage.
- [28] D.V. Nanopoulos (1995) *Theory of Brain function, Quantum Mechanics, and Superstrings*, CERN-TH/95-128, hep-ph/9505374.
- [29] J. Hitt (1999), *This is Your Brain on God*, Wired Archive,
http://www.wired.com/wired/archive/7.11/persinger_pr.html.
- [30] P. Pietch, (1972), *Shuffle Brain: the The Quest for Hologramic Mind*,
<http://www.indiana.edu/~pietsch/shufflebrain-book00.html>.

- [31] S. Tippit (2001), *Mind continues after brain dies, scientist says*. An article of Guardian about the work of University of Southampton research team about NDE.
<http://www.sun-sentinel.com/news/local/southflorida/sfl-628brain>.
- [32] Paramahansa Yogananda (1993) ,*Autobiography of Yogi*. Self-Realization Fellowship.
- [33] M. Pines, *A Secret Sense in The Human Nose?*
<http://www.hhmi.org/senses/>.
- [34] Fröhlich, H. (1975) *The extraordinary dielectric properties of biological materials and the action of enzymes*, Proc. Natl. Acad. Sci. 72:4211-4215.
- [35] Albrecht-Buehler,G. (1991) *Surface extensions of 3T3 cells towards distant infrared light sources*. Journal of Cell Biology, 114, 493-502.
- [36] G. Pollack (200?), *Cells, Gels and the Engines of Life*, Ebner and Sons.
<http://www.cellsandgels.com/> .
- [37] Koruga, D. (1974),*Microtubule screw symmetry: packing of spheres as a latent bioinformation code*. Ann. Ny. Acad. Sci. 466, 953-955.

Effects of em fields on living matter

- [38] R. O. Becker and G. Selden (1990) *The Body Electric: Electromagnetism and the Foundation of Life*. William Morrow & Company, Inc., New York.
- [39] J. Benveniste *et al* (1988). *Human basophil degranulation triggered by very dilute antiserum against IgE*. Nature 333:816-818.
- [40] J. Benveniste *et al* (198?). *Transatlantic transfer of digitized antigen signal by telephone link*. Journal of Allergy and Clinical Immunology. 99:S175 (abs.).FFor recent work about digital biology and further references about the work of Benveniste and collaborators see <http://www.digibio.com/> .
- [41] C. F. Blackman (1994), "Effect of Electrical and Magnetic Fields on the Nervous System" in *The Vulnerable Brain and Enviromental Risks, Vol. 3, Toxins in Air and Water* (eds. R. L. Isaacson and K. F. Jensen). Plenum Press, New York, pp. 331-355.
- [42] M. Buzzacchi, E. Del Giudice and G. Preparata (1998), *Sonoluminescence unveiled?*, <http://xxx.lanl.gov/quant-ph/9804006> .
- [43] Callahan, P. S. (1977). *Moth and Candle: the Candle Flame as a Sexual Mimic of the Coded Infrared Wavelengths from a Moth Sex Scent*. Applied Optics. 16(12) 3089-3097.
- [44] N. Cherry (2000), Conference report on effects of ELF fields on brain, <http://www.tassie.net.au/emfacts/icnirp.txt> .
- [45] P. Gariaev *et al* (2000), *The DNA-wave biocomputer*, International Journal of Computing Anticipatory Systems. Ed. Daniel Dubois, Published by CHAOS, Vol. 10, 2001.
- [46] P. Marcer and W. Schempp (1996), *A Mathematically Specified Template For DNA and the Genetic Code in Terms of Physical Process of Quantum Holography*. Proceedings of the Greenwich Symposium of Living Computers, eds. A. Fedorec and P. Marver. 45-62.
P. Marcer and W. Schempp (1998), *The brain as a conscious system*, International Journal of General Systems, 27, 1/3, 231,248.

- [47] P. Marcer *et al* (2000), *Quantum Millennium, Quantum Universe, Quantum Biosphere, Quantum Man- or What Physicists can teach Biologists, and Biology, Physics*, International Journal of Computing Anticipatory Systems. Ed. Daniel Dubois, Published by CHAOS, Vol. 10, 2001.
- [48] P. Gariaev and A. Junin (1989), *Energy*, no 10,45-62 v. 11 (in Russian).
- [49] Jibu,M., Hagan,S., Hameroff, S.R., Pribram,K.H., Yasue,K. (1994). *Quantum optical coherence in cytoskeletal microtubules: implications for brain function*. *Biosystems*,32, 195-209.
- [50] Abstracts of talks 'Macroscopic model of collective microtubule instability: supercatastrophes' and 'Macroscopic model of elementary act of perception and braining' held in Tucson II. conference:<http://www.eu.arizona.edu/~uaextaed/conferen/consc.html>.
- [51] L. Milgrom (2001), *Thanks for the memory*. An article in Guardian about the work of professor M. Ennis of Queen's University Belfast supporting the observations of Dr. J. Benveniste about water memory.
<http://www.guardian.co.uk/Archive/Article/0,4273,4152521,00.html> .
- [52] W. Nagl, M. Rattemayer and F.A. Popp (1981), *Evidence of Photon Emission from DNA in Living Systems*, in *Naturwissenschaften*, Vol. 68, No 5, 577.
- [53] F. A. Popp (2000), *Some features of biophotons and their interpretation in terms of coherent states*. *Biophotonics and Coherent Systems*. Proc. 2nd A. Gurwitsch Conference and Additional Contributions. Moscow University Press. Ed. I. Belousov *et al*, 117-133.
- [54] C. Smith (2001), *Learning From Water, A Possible Quantum Computing Medium*, talk in CASYS'2001, 5th international conference on Computing Anticipating Systems held in Liege, Belgium, August 13-18. Abstract book published by Chaos.
- [55] D. Yarrow (1990), *Spin the tale of the dragon*, review article on biomagnetism, <http://www.ratical.org/ratville/RofD2.html>.

References related to anomalies

- [56] Marc Bean , *Thomas Townsend Brown*,
<http://www.virtualtheater.com/ttbrown-01.html> .
- [57] Gaston Burrige (1994), *Townsend Brown and his antigravity discs*.
<http://www.spiritweb.com/Spirit/anti-gravity.html> .
- [58] Mason Rose (1952) , "*The Flying Saucer*". *The Application of the Biefeld-Brown Effect to the Solution of the Problems of Space Navigation*. Published in *Science and Invention*, August 1929.
<http://www.soteria.com/brown/docs/rose.htm>.